

Weak- L^1 estimates and ergodic theorems

Ciprian Demeter and Anthony Quas

ABSTRACT. We prove that for any dynamical system (X, Σ, m, T) , the maximal operator defined by

$$N^* f(x) = \sup_n \frac{1}{n} \# \left\{ 1 \leq i : \frac{f(T^i x)}{i} \geq \frac{1}{n} \right\}$$

is almost everywhere finite for f in the Orlicz class $L \log \log L(X)$, extending a result of Assani [2]. As an application, a weighted return times theorem is also proved.

CONTENTS

1. Introduction		169
2. Main results		171
References		174

1. Introduction

Let T be a measure preserving transformation of a probability space (X, Σ, m) . We call (X, Σ, m, T) a dynamical system. The following return times theorem was proved in [4]:

Theorem 1 (Bourgain). *Let $1 \leq p \leq \infty$ and let $1/p + 1/q = 1$. For each dynamical system (X, Σ, m, T) and $f \in L^p(X)$, there is a set $X_0 \subset X$ of full measure, such that for any other dynamical system (Y, \mathcal{F}, μ, S) , $g \in L^q(Y)$ and $x \in X_0$, the limit,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(T^k x) g(S^k y),$$

exists for μ a.e. y .

One of the most interesting unanswered questions that emerges from this result is whether or not the fact that f and g lie in dual spaces is in general necessary in order to have a positive result. Neither of the existing proofs of Theorem 1 gives any indication on this, since each of them relies on Hölder’s inequality.

Received August 14, 2003.

Mathematics Subject Classification. 37A30, 46E30, 60F15.

Key words and phrases. Return times theorem, Orlicz spaces.

The second author’s research was partially supported by NSF Grant DMS-0200703.

On the other hand, if (gS^k) is replaced with a sequence (ξ_k) of independent identically distributed random variables such that $\mathbb{E}(|\xi_1|) < \infty$, then the following criterion of B. Jamison, S. Orey and W. Pruitt [5] proves to be an excellent tool to break the duality.

Theorem 2 (Jamison, Orey and Pruitt). *Let (a_k) be a sequence of positive real numbers and let $N^* = \sup_n \frac{1}{n} \#\{k : a_k / \sum_{i=1}^k a_i \geq 1/n\}$, then the following are equivalent:*

1. $N^* < \infty$.
2. For any i.i.d. sequence of random variables (ξ_k) such that $\mathbb{E}(|\xi_1|) < \infty$, defining a new sequence (Ξ_n) of random variables by

$$\Xi_n(\omega) = \frac{\sum_{k=1}^n a_k \xi_k(\omega)}{\sum_{k=1}^n a_k},$$

the sequence (Ξ_n) converges pointwise almost surely.

Motivated by this criterion, Assani [1] introduced the following maximal function: given $f \in L^1(X)$, consider

$$N^* f(x) = \sup_n \frac{1}{n} \#\left\{1 \leq i : \frac{f(T^i x)}{i} \geq \frac{1}{n}\right\}.$$

He proved in [2] for $f \in L \log L(X)$, $N^* f \in L^1$ and in particular $N^* f(x) < \infty$ for a.e. x . Based on this and Theorem 2, the following “duality-breaking” version of Theorem 1 follows almost immediately:

Corollary 3 (Assani). *Let (X, Σ, m, T) be a measure-preserving transformation and let the function f satisfy $\int |f| \log^+ |f| dm < \infty$, that is $f \in L \log L(X)$. Then there is a set $X_0 \subset X$ of full measure, such that for any sequence (ξ_k) of i.i.d. random variables on the probability space $(\Omega, \mathcal{F}, \mu)$ with $\xi_1 \in L^1(\Omega)$ and any $x \in X_0$*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(T^k x) \xi_k(\omega)$$

exists for μ a.e. ω .

Moreover in [1] it is proved that if Theorem 1 is true for $p = q = 1$, then $N^* f(x)$ must be finite almost everywhere for all $f \in L^1(X)$. This connection sheds more light on the importance of the operator N^* and motivates its further study.

In the next section we will prove the finiteness of N^* for functions in the larger class $L \log \log L$. Note that while Assani shows that $N^* f \in L^1$ for $f \in L \log L$, our result establishes that $N^* f \in L^{1,\infty}$ for $f \in L \log \log L$ (i.e., that $\sup_t t m\{x : N^* f(x) > t\} < \infty$) so that while our hypothesis is weaker, so is our conclusion. Note however that since our conclusion implies that $N^* f(x) < \infty$ for almost every x , it is sufficient to imply a corollary like Corollary 3 in the case where $f \in L \log \log L$.

In a preprint that appeared at around the time this paper was submitted, Assani, Buczolic, and Mauldin [3] show that there exists an $f \in L^1(X)$ such that $N^* f(x) = \infty$ almost everywhere.

2. Main results

Throughout this section we will denote the natural logarithm of x by $\log x$ and the weak- L^1 norm of f by

$$\|f\|_{1,\infty} = \sup_{\lambda>0} \lambda m\{x : |f(x)| > \lambda\}.$$

We will also need to refer to the *entropy* of a sequence of positive real numbers. Specifically, for a sequence (a_n) of nonnegative real numbers (not all 0), define the entropy by

$$H((a_n)) = \sum_n -\frac{a_n}{\sum_j a_j} \log \left(\frac{a_n}{\sum_j a_j} \right),$$

under the convention $0 \log 0 = 0$.

We define f^* to be the ergodic maximal function, $f^*(x) = \sup_n \left| \frac{1}{n} \sum_{k=1}^n f(T^k x) \right|$. The maximal ergodic theorem asserts that $\|f^*\|_{1,\infty} \leq \|f\|_1$ for all $f \in L^1(X)$. The following inequality from [7] turns out to be extremely useful to our investigation:

Lemma 4. *Suppose that for $i = 1, 2, \dots$, $g_i(x)$ is an $L_{1,\infty}$ function on a measure space such that $\sum \|g_i\|_{1,\infty} < \infty$. Then*

$$\left\| \sum_{i=1}^{\infty} g_i \right\|_{1,\infty} \leq 2(K+2) \sum_{i=1}^{\infty} \|g_i\|_{1,\infty},$$

where K is the entropy of the sequence $(\|g_n\|_{1,\infty})$.

We can now prove our main result.

Theorem 5. *For each dynamical system (X, Σ, m, T) and each $f \in L \log \log L(X)$ (that is f satisfying $\int |f| \log^+ \log^+ |f| dm < \infty$), $N^* f(x) < \infty$ for a.e. x .*

Proof. It is enough to consider f positive. Making use of the fact that $f(x) \leq \sum_{i=1}^{\infty} 2^i \chi_{A_i}(x)$, where $A_i = \{x : 2^{i-1} < f(x) \leq 2^i\}$ for $i \geq 2$ and $A_1 = \{x : f(x) \leq 2\}$, it easily follows that for each n ,

$$\frac{1}{n} \# \left\{ k \geq 1 : \frac{f(T^k x)}{k} \geq 1/n \right\} \leq \frac{1}{n} \sum_{i=1}^{\infty} \sum_{k=1}^{n2^i} \chi_{A_i}(T^k x) \leq \sum_{i=1}^{\infty} 2^i (\chi_{A_i})^*(x).$$

We will show that the last term in the above inequality is finite a.e. by proving that its $L_{1,\infty}$ norm is finite.

Let $a_i = 2^i \|\chi_{A_i}\|_{1,\infty}$. By the maximal ergodic theorem, we see that $a_i \leq 2^i m(A_i)$. The fact that $f \in L \log \log L$ implies that $\sum_i a_i \log i < \infty$ (and hence clearly M , which we define to be $\sum_i a_i$, is finite). From the lemma above, we see that it is sufficient to show that the entropy of the sequence (a_i) is finite: $\sum_i -a_i/M \log(a_i/M) < \infty$. One quickly sees that this is equivalent to establishing $\sum_i -a_i \log a_i < \infty$.

Consider now $S_1 = \{i : a_i \leq 1/i^2\}$ and $S_2 = \{i : a_i > 1/i^2\}$. Now

$$\sum_{i \in S_1} -a_i \log a_i \leq 1 + \sum_{j=2}^{\infty} \log(j^2)/j^2 < \infty$$

since $\psi(t) = -t \log t$ is increasing on $[0, 1/e]$. On the other hand

$$\sum_{i \in S_2} -a_i \log a_i < 2 \sum_{i \in S_2} a_i \log i < \infty.$$

□

Corollary 6. *For each dynamical system (X, Σ, m, T) and nonnegative function $f \in L \log \log L(X)$, there is a set $X_0 \subset X$ of full measure, such that for any sequence (ξ_k) of i.i.d. random variables on the probability space $(\Omega, \mathcal{F}, \mu)$ with $\xi_1 \in L^1(\Omega)$ and any $x \in X_0$*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(T^k x) \xi_k(\omega)$$

exists for μ a.e. ω .

There does not seem to be a better way of exploiting Lemma 4 in order to extend even more the class of functions for which N^*f is almost everywhere finite. Moreover, as we show in the following proposition, the inequality in Lemma 4 is sharp up to a constant. We note that a more general version of this proposition appears in work of Kalton [6].

Proposition 7. *Given positive numbers a_1, \dots, a_n , there exist functions g_1, \dots, g_n with $\|g_i\|_{1,\infty} = a_i$ such that $\|g_1 + \dots + g_n\|_{1,\infty} \geq \frac{1}{6}(2+K) \sum \|g_i\|_{1,\infty}$, where K is the entropy of the sequence (a_i) .*

Proof. For each i , let ξ_i be a random variable taking the value $1/n$ with probability $(1 - a_i)^{n-1} a_i$. Moreover, the ξ_i 's will be chosen to be independent. One can then check that $\mathbb{P}(\xi_i > \lambda) \leq a_i/\lambda$ while $\mathbb{P}(\xi_i \geq 1 - \epsilon) = a_i$ for ϵ small enough, so that $\|\xi_i\|_{1,\infty} = a_i$.

We see that

$$\begin{aligned} \mathbb{E}(\xi_i) &= \sum_{n=1}^{\infty} a_i \frac{(1 - a_i)^{n-1}}{n} \\ &= -\frac{a_i}{1 - a_i} \log a_i \geq -a_i \log a_i. \end{aligned}$$

Similarly, we see that

$$\mathbb{E}(\xi_i^2) = a_i \sum_{n=1}^{\infty} \frac{1}{n^2} (1 - a_i)^{n-1} \leq a_i \sum_{n=1}^{\infty} \frac{1}{n^2} \leq 2a_i.$$

In particular, setting $\Xi = \xi_1 + \dots + \xi_n$, we see that $\mathbb{E}(\Xi) \geq K$ but $\text{Var}(\Xi) \leq 2$.

Using Tchebychev's inequality, we see that

$$\mathbb{P}(\Xi \geq K - 2) \geq \mathbb{P}(|\Xi - \mathbb{E}(\Xi)| \leq 2) \geq 1 - \frac{\text{Var}(\Xi)}{2^2} \geq \frac{1}{2}.$$

If $K > 4$, we have $\mathbb{P}(\Xi \geq K/2) \geq \frac{1}{2}$ so that the weak- L^1 norm exceeds $K/4$, which in turn exceeds $(K+2)/6$. If $K \leq 4$, take f to be any function of weak L^1 norm 1 and let $f_n = a_n f$, so that $\|f_n\|_{1,\infty} = a_n$. Then $\sum f_i = f$, so that $\|\sum f_i\|_{1,\infty} = 1 \geq \frac{1}{6}(K+2) \sum \|f_i\|_{1,\infty}$. This completes the proof of the proposition. □

Remark 8. Note that although $f \in L \log \log L$ is sufficient to guarantee that $N^*f < \infty$ almost everywhere, there are functions f outside $L \log \log L(X)$, for which $N^*f(x) < \infty$ for a.e. x . In particular, it is easy to construct functions outside $L \log \log L$ for which the entropy computed in Theorem 5 is finite, guaranteeing the finiteness of N^*f .

Further, if we are willing to restrict the system, we see that no condition on the distribution of f ensures the divergence of $N^*f(x)$. Specifically, Lemma 1 of [1] guarantees that whenever $T^k f$ are independent (identically distributed) random variables with an arbitrary L^1 distribution, then $N^*f(x) < \infty$ for a.e. x .

Another consequence of Theorem 5 is the following weighted version of Corollary 3.

Theorem 9. *For each dynamical system (X, Σ, m, T) and $f \in L^1(X)$, there is a set $X_0 \subset X$ of full measure, such that for any sequence (ξ_k) of i.i.d. random variables on the probability space $(\Omega, \mathcal{F}, \mu)$ with $\xi_1 \in L^1(\Omega)$ and any $x \in X_0$*

$$\lim_{n \rightarrow \infty} \frac{1}{n \log \log n} \sum_{k=1}^n f(T^k x) \xi_k(\omega) = 0$$

for μ a.e. ω .

The proof will be based on the following relative of Theorem 5. Define

$$L^*f(x) = \sup_n \frac{1}{n} \# \left\{ 1 \leq i : \frac{f(T^i x)}{i \log \log i} \geq \frac{1}{n} \right\}.$$

Lemma 10. *Let (X, Σ, m, T) be a measure-preserving system. For each $f \in L_1(X)$, $L^*f(x) < \infty$ for a.e. x .*

Proof. As usual, we can assume f is positive. Fix an $n \in \mathbb{N}$. Using the fact that $f(x) \leq \sum_{i=1}^{\infty} 2^i \chi_{A_i}(x)$ we get that

$$\frac{1}{n} \# \left\{ 1 \leq k : \frac{f(T^k x)}{k \log \log k} \geq 1/n \right\} \leq \frac{1}{n} \sum_{i=1}^{\infty} \sum_{k=1}^{p_i} \chi_{A_i}(T^k x) \leq \frac{1}{n} \sum_{i=1}^{\infty} p_i (\chi_{A_i})^*(x)$$

where p_i is the largest integer such that $p_i(\log \log p_i) \leq n2^i$. Letting $\phi: (1, \infty) \rightarrow \mathbb{R}$ be the increasing function $\phi(x) = x \log \log x$, we see that $p_i \leq \phi^{-1}(n2^i)$. We claim that there exists a $C > 0$ such that $p_i \leq C \frac{2^i n}{\log(i+1)}$ for all $i, n \in \mathbb{N}$. To see this, we check the existence of a C such that $\phi^{-1}(2^x) \leq C \frac{2^x}{\log(x+1)}$ or equivalently $2^x \leq \phi(C \frac{2^x}{\log(x+1)})$ for all $x \geq 0$. Hence

$$\sup_n \frac{1}{n} \# \left\{ 1 \leq i : \frac{f(T^i x)}{i \log \log i} \geq \frac{1}{n} \right\} \leq C \sum_{i=1}^{\infty} \left(\frac{2^i}{\log(i+1)} \right) (\chi_{A_i})^*(x).$$

Based on Lemma 4 and on the maximal ergodic theorem, it suffices to prove that $\sum_{i=1}^{\infty} \left(\frac{2^i \|\chi_{A_i}\|_1}{\log(i+1)} \right) \log \left(\frac{2^i \|\chi_{A_i}\|_1}{\log(i+1)} \right) < \infty$. By splitting the sum in two parts depending on whether or not $\frac{2^i \|\chi_{A_i}\|_1}{\log(i+1)} < \frac{1}{i^2}$ and reasoning as in the proof of Theorem 5, it easily follows that the sum from above is finite. \square

Proof of Theorem 9. It suffices to assume that both f and ξ_1 are positive. According to the previous lemma, let X_0 the subset of full measure of X containing all the points x for which $L^*f(x) < \infty$. For a fixed $x \in X_0$ denote $w_k := f(T^k x)$ and also $W_k := k \log \log k$. The argument of Jamison, Orey and Pruitt from [5] can be extended with really no essential changes to this case, to conclude that since

$$\sup_n \frac{1}{n} \# \left\{ 1 \leq i : \frac{w_i}{W_i} \geq \frac{1}{n} \right\} < \infty,$$

$$\lim_{n \rightarrow \infty} \frac{1}{W_n} \sum_{k=1}^n w_k \xi_k(\omega) = 0$$

for μ a.e. ω . □

Remark 11. It is not known whether in Theorem 9 the weight $n \log \log n$ can be replaced with a smaller one, like $n \log \log \log n$. Any improvement on this weight will necessarily have behind it an extension of the result of Theorem 5 to a larger Orlicz class. On the other hand, a combination of Theorem 2 and the result from [3], shows that this weight can not be chosen to be n .

Remark 12. It would be interesting to find the largest Orlicz class that would guarantee that $N^*f(x) < \infty$ almost everywhere. The above establishes that such an Orlicz class would contain $L \log \log L$.

A careful examination of the proof of [3] demonstrates that in any Orlicz class with an essentially smaller weight than the class $L \log \log \log L$, there exists a function f such that $N^*f(x) = \infty$ almost everywhere.

In particular, these two results demonstrate that the largest Orlicz class that would guarantee that $N^*f(x) < \infty$ almost everywhere lies between $L \log \log L$ and $L \log \log \log L$.

References

- [1] I. Assani, *Strong laws for weighted sums of independent identically distributed random variables*, Duke Math. J. **88** (1997), 217–246, MR 1455518 (98g:60059), Zbl 0883.60023.
- [2] I. Assani, *Convergence of the p -Series for stationary sequences*, New York J. Math. **3A** (1997), 15–30, MR 1604577 (99c:60061), Zbl 0915.28009.
- [3] I. Assani, Z. Buczolich, and R. Mauldin, *An L^1 counting problem in ergodic theory*, arXiv math.DS/0307384.
- [4] J. Bourgain, H. Furstenberg, Y. Katznelson and D. Ornstein, *Return times of dynamical systems* (appendix to Bourgain’s ‘Pointwise ergodic theorems for arithmetic sets’), Inst. Hautes Études Sci. Publ. Math. **69** (1989), 47–50, MR 1019960 (90k:28030), Zbl 0705.28008.
- [5] B. Jamison, S. Orey, and W. Pruitt, *Convergence of weighted averages of independent random variables*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **4** (1965), 40–44, MR 0182044 (31 #6268), Zbl 0141.16404.
- [6] N. J. Kalton, *Sequences of random variables in L_p for $p < 1$* , J. Reine Angew. Math. **329** (1981), 204–214, MR 0636454 (83a:46040), Zbl 0461.60020.
- [7] E. M. Stein and N. J. Weiss, *On the convergence of Poisson integrals*, Trans. Amer. Math. Soc. **140** (1969), 35–54, MR 0241685 (39 #3024), Zbl 0182.10801.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA, URBANA, IL 61801
demeter@math.uiuc.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF MEMPHIS, MEMPHIS, TN 38152-6429
aquas@memphis.edu

This paper is available via <http://nyjm.albany.edu:8000/j/2004/10-10.html>.