

# Regular AF subalgebras of some crossed products

Catalin Rada

ABSTRACT. Let  $C(X) \rtimes_T \mathbb{Z}$  be the crossed product associated to a dynamical system  $(X, T)$ . We characterize the regular AF subalgebras of  $C(X) \rtimes_T \mathbb{Z}$  that can arise as the algebra  $A_Y = \langle C(X), uC_0(X \setminus Y) \rangle$  for some closed subset  $Y$  of  $X$ . We also characterize the minimal homeomorphisms in  $A_Y$  terms.

## CONTENTS

1. Regular AF subalgebras	431
2. Minimality	433
References	434

## 1. Regular AF subalgebras

Let  $X$  be the Cantor set and let  $T : X \rightarrow X$  be a minimal homeomorphism. Denote by  $u$  the canonical unitary implementing the action  $\mathbb{Z} \ni n \rightarrow \alpha_T^n \in \text{Aut}(C(X))$ , where  $\alpha_T : C(X) \rightarrow C(X)$  is given by  $\alpha_T(f) = f \circ T^{-1}$  for every  $f \in C(X)$  and by  $C(X) \rtimes_T \mathbb{Z}$  the crossed product  $C(X) \rtimes_{\alpha_T} \mathbb{Z}$ . Set  $C_0(X \setminus Y) = \{f \in C(X) \mid f|_Y = 0\}$ . Recall the following theorem of Putnam ([2]):

**Theorem 1.1.** *For any nonempty closed subset  $Y$  of  $X$ , the  $C^*$ -subalgebra of  $C(X) \rtimes_T \mathbb{Z}$  generated by  $C(X)$  and  $uC_0(X \setminus Y)$  is an AF algebra.*

Set  $A_Y = \langle C(X), uC_0(X \setminus Y) \rangle$ . For any  $C^*$ -subalgebra  $A$  of  $C(X) \rtimes_T \mathbb{Z}$  containing  $C(X)$ , the normalizer of  $C(X)$  in  $U(A)$  is given by  $N(C(X), A) = \{v \in U(A) \mid vC(X)v^* = C(X)\}$ . (Here  $U(A)$  is the set of unitary elements in  $A$ .) The algebra  $A$  is called *regular* if  $N(C(X), A)$  generates  $A$ . It is

---

Received August 21, 2009.

*Mathematics Subject Classification.* Primary 46L05; Secondary 37B05.

*Key words and phrases.* Crossed products,  $C^*$ -algebras.

well-known that  $A_Y$  is regular. Another property of  $A_Y$  is contained in the following result of Poon (see [1]):

**Theorem 1.2.** *Let  $Y$  be a nonempty subset of  $X$ . For every  $n \geq 1$  and any clopen subset  $W$  of  $X$  one has that  $u^n \chi_W (= u \chi_{T^{n-1}(W)} u \chi_{T^{n-2}(W)} \dots u \chi_W) \in A_Y$  if and only if  $u \chi_{T^{n-1}(W)}, u \chi_{T^{n-2}(W)}, \dots, u \chi_W \in A_Y$ . (Here  $\chi_W$  denotes the characteristic function of  $W$ .)*

We say that a  $C^*$ -subalgebra  $A$  of the crossed product  $C(X) \rtimes_T \mathbb{Z}$  has property  $\star$  if for any  $n \geq 1$  and any clopen subset  $W$  of  $X$ ,

$$u \chi_{T^{n-1}(W)}, u \chi_{T^{n-2}(W)}, \dots, u \chi_W \in A$$

whenever  $u^n \chi_W \in A$ .

**Theorem 1.3.** *For any regular AF subalgebra  $A$  of  $C(X) \rtimes_T \mathbb{Z}$  that contains  $C(X)$  and has property  $\star$ , there is a closed nonempty set  $Y$  of  $X$  such that  $A_Y = A$ .*

Let us find the natural candidate for  $Y$ . Consider the set

$$I = \{f \in C(X) \mid uf \in A\} \subseteq C(X).$$

**Lemma 1.4.** *The subset  $I$  is a closed ideal of  $C(X)$ .*

**Proof.** If  $f, g \in I$  then  $uf, ug \in A$ , and since  $A$  is an algebra  $u(f + g) = uf + ug \in A$ , so  $f + g \in I$ . If  $f \in I$  and  $g \in C(X)$  then  $uf \in A$  and since  $g \in C(X) \subseteq A$  we get  $ufg \in A$ , hence  $fg \in I$ . Remark that since  $g \circ T^{-1} \in C(X) \subseteq A$  and  $uf \in A$  we get that  $ugf = (g \circ T^{-1})uf \in A$ , hence  $gf \in I$ . Suppose that  $f_n \rightarrow f$  and  $f_n \in I$ . Then  $uf_n \rightarrow uf$ , and since  $uf_n \in A$  it follows that  $uf \in A$ , hence  $f \in I$ .  $\square$

Since  $I$  is an ideal in  $C(X)$  we know that  $I = C_0(X \setminus Y)$  for some closed subset  $Y$  of  $X$ . Since  $A$  is AF we see that  $u$  is not in  $A$ , so  $Y$  is not empty.

**Lemma 1.5.** *The inclusion  $A_Y \subseteq A$  is always valid.*

**Proof.** It is sufficient to show that the generators of  $A_Y$  are in  $A$ . By the definition of  $A$  we know that  $C(X) \subseteq A$ . If  $f \in C_0(X \setminus Y) = I$  then  $uf \in A$ , hence  $uC_0(X \setminus Y) \subseteq A$ . Therefore  $A_Y \subseteq A$ .  $\square$

**Lemma 1.6.** *The inclusion  $A \subseteq A_Y$  holds.*

**Proof.** Since  $A$  is a regular algebra it is enough to show that each unitary in  $N(C(X), A)$  is also in  $A_Y$ . Letting  $v$  be a unitary in  $N(C(X), A)$ , it is known from Lemma 5.1 of [2] that  $v = f \sum_{n \in \mathbb{Z}} p_n u^n$ , where  $f$  is a unitary in  $C(X)$ , every  $p_n$  is a projection in  $C(X)$ , only finitely many  $p_n$  are different than 0,  $p_n p_k = 0$  for  $n \neq k$  and  $\sum_{n \in \mathbb{Z}} p_n = 1$ . Since the projections  $p_n$  are orthogonal, we obtain that  $p_n u^n \in A$  for every  $n \in \mathbb{Z}$ .

We show that  $p_n u^n \in A_Y$ . Suppose that  $n \geq 1$ . If  $p_n = \chi_W$  for a clopen subset  $W$  of  $X$ , from the chain of equalities  $p_n u^n = u^n (p_n \circ T^n) = u^n \chi_{T^{-n}(W)} = u \chi_{T^{-1}(W)} u \chi_{T^{-2}(W)} \dots u \chi_{T^{-n}(W)}$  and the  $\star$  condition,

$$u \chi_{T^{-1}(W)}, u \chi_{T^{-2}(W)}, \dots, u \chi_{T^{-n}(W)} \in A.$$

Hence we see that  $\chi_{T^{-1}(W)}, \chi_{T^{-2}(W)}, \dots, \chi_{T^{-n}(W)} \in I = C_0(X \setminus Y)$ . From the decomposition  $p_n u^n = u \chi_{T^{-1}(W)} u \chi_{T^{-2}(W)} \dots u \chi_{T^{-n}(W)}$  conclude that  $p_n u^n \in A_Y$ . If  $p_{-n} u^{-n} \in A$  for some  $n \geq 1$ , we note that  $u^n p_{-n} = (p_{-n} u^{-n})^* \in A$ , and as above one gets that  $u^n p_{-n} \in A_Y$ , and therefore  $p_{-n} u^{-n} \in A_Y$ .  $\square$

**Remark 1.7.** (1) Condition  $\star$  can not be removed. To this end, let  $T$  be a minimal homeomorphism of  $X$  such that  $T^2$  is also minimal. We embed  $C(X) \rtimes_{T^2} \mathbb{Z}$  into  $C(X) \rtimes_T \mathbb{Z}$  via  $C(X) \ni f \rightarrow f \in C(X)$  and  $v \rightarrow u^2$ , where  $v$  is the canonical unitary in  $C(X) \rtimes_{T^2} \mathbb{Z}$  implementing the action generated by  $T^2$ . Let  $Y$  be a closed, nonvoid subset of  $X$ , with  $Y \neq X$ , and consider the regular AF subalgebra  $\langle C(X), v C_0(X \setminus Y) \rangle$  of  $C(X) \rtimes_{T^2} \mathbb{Z}$ . We can see this subalgebra as  $\langle C(X), u^2 C_0(X \setminus Y) \rangle$  in the crossed product  $C(X) \rtimes_T \mathbb{Z}$ . The last algebra is then again regular and AF, not satisfying the condition  $\star$  and not of the form  $\langle C(X), u C_0(X \setminus Z) \rangle$  for any nonempty closed subset  $Z$  of  $X$ .

(2) If  $\star$  is removed and  $Y$  is a singleton it was noted by Poon (see Corollary 4.4 of [1]) that Lemma 1.6 still holds.

## 2. Minimality

In this section we give a characterization of the minimal homeomorphisms in terms of approximately finitely-dimensionality of  $A_Y$ s. If  $X$  is a zero-dimensional compact space and  $T$  a self-homeomorphism of  $X$ , not necessarily minimal, then we can still define the  $C^*$ -subalgebra  $\langle C(X), u C_0(X \setminus Y) \rangle$ . It was proved in [1] that  $A_Y$  is an AF algebra iff  $X = \bigcup_{n \in \mathbb{Z}} T^n(W)$  for every clopen subset  $W$  containing  $Y$ .

**Proposition 2.1.** *Let  $X$  be a zero-dimensional compact space and  $T : X \rightarrow X$  a self-homeomorphism. Then  $T$  is minimal iff  $A_Y$  is AF for any closed nonempty subset  $Y$  of  $X$ .*

**Proof.** Suppose that  $A_Y$  is AF for any nonempty closed subset  $Y$  of  $X$ . If there is a proper closed subset  $Y$  of  $X$  such that  $T(Y) = Y$  (i.e.,  $Y \neq \emptyset, Y \neq X$ ) then  $Y$  is not clopen. Pick an element  $x_0 \in X \setminus Y$  and note again that  $Y \cup \{x_0\} \neq X$  (otherwise  $Y$  were clopen). Let  $W$  be a clopen subset of  $X$  such that  $Y \cup \{x_0\} \subseteq W \subseteq X$  and  $Y \cup \{x_0\} \neq W, W \neq X$ .

Since  $x_0$  is not in  $Y$  and  $x_0 \in W$  there is a clopen subset  $U$  such that  $x_0 \in U \subseteq W$  and  $U \cap Y = \emptyset$ . But  $W^c \cup \{x_0\}$  is a closed subset of  $X$  that is included in the clopen subset  $W^c \cup U$ . By Poon's result (cited above) one gets that

$X = \bigcup_{n \in \mathbb{Z}} T^n(W^c \cup U)$ . Since  $Y$  is not empty let  $y_0 \in Y$ , we get that  $y_0 \in T^{n_0}(W^c \cup U)$  for some  $n_0 \in \mathbb{Z}$ . We distinguish two cases. If  $y_0 \in T^{n_0}(W^c)$ , taking into account that also  $y_0 \in Y$  implies that  $y_0 \in T^{n_0}(Y)$ , one gets that  $y_0 \in T^{n_0}(W^c) \cap T^{n_0}(Y) = T^{n_0}(W^c \cap Y) = T^{n_0}(\emptyset) = \emptyset$ , a contradiction. If  $y_0 \in T^{n_0}(U)$  then again  $y_0 \in T^{n_0}(U) \cap T^{n_0}(Y) = T^{n_0}(U \cap Y) = T^{n_0}(\emptyset) = \emptyset$ , contradiction.

The other implication is always true by [2]. □

**Acknowledgements.** We are deeply indebted to Professor David Handelman for many useful discussions and continuous support.

## References

- [1] POON, YIU TUNG. AF subalgebras of certain crossed products. *Rocky Mountain J. Math.* **20** (1990) 527–537. MR1065849 (91k:46078), Zbl 0727.46044.
- [2] PUTNAM, IAN F. The  $C^*$ -algebras associated with minimal homeomorphisms of the Cantor set. *Pacific J. Math.* **136** (1989) 329–353. MR0978619 (90a:46184), Zbl 0631.46068.

UNIVERSITY OF OTTAWA, DEPARTMENT OF MATHEMATICS AND STATISTICS, OTTAWA, ON, CANADA  
 catalin.rada@alumni.uottawa.ca

This paper is available via <http://nyjm.albany.edu/j/2009/15-24.html>.