

Concordances from connected sums of torus knots to L -space knots

Charles Livingston

ABSTRACT. If a knot is a nontrivial connected sum of positive torus knots, then it is not concordant to an L -space knot.

CONTENTS

1. Introduction	233
2. Torus knot Alexander polynomials and signature functions	234
2.1. Alexander polynomials of torus knots	234
2.2. Signature functions of torus knots	235
3. Concordances to L -space knots	237
4. Generalizations	237
References	238

1. Introduction

In a recent paper about involutive knot Floer homology, Zemke [19] observed that invariants arising from Heegaard Floer knot homology can obstruct a knot from being concordant to an L -space knot. He considered examples of connected sums of torus knots, such as $T(4, 5) \# T(4, 5)$ and $-T(3, 4) \# -T(4, 5) \# T(5, 6)$, and noted that in some cases alternative obstructions are available. Here we use classical knot invariants along with the Ozsváth–Szabó tau invariant [14], $\tau(K)$, to prove the following theorem.

Theorem. *Let $\{(p_i, q_i)\}_{i=1, \dots, n}$ be a set of pairs of relatively prime positive integers with $2 \leq p_i < q_i$ for all i and with $n > 1$. Then $\#_i T(p_i, q_i)$ is not concordant to an L -space knot.*

The main idea of the proof can be illustrated with the example $K = T(4, 5) \# T(4, 5)$. Since $\tau(K) = 12$, if K is concordant to a knot J , then $\tau(J) = 12$. If J is an L -space knot, then the Alexander polynomial of J

Received January 15, 2018.

2010 *Mathematics Subject Classification.* 57M25.

Key words and phrases. Knot concordance, L -space knot, torus knot.

The author was supported by NSF-DMS-1505586.

is of degree 24. The Alexander polynomial $\Delta_K(t)$ is the product of cyclotomic polynomials $\phi_{10}(t)^2\phi_{20}(t)^2$. Since the Levine–Tristram signature function [10, 18] for K jumps by four at the roots of $\phi_{20}(t)$ and $\phi_{10}(t)$, the same is true for J . This implies that $\phi_{10}(t)^2\phi_{20}(t)^2$ divides $\Delta_J(t)$. By degree considerations, this implies $\Delta_J(t) = \phi_{10}^2(t)\phi_{20}^2(t) = t^{24} - 2t^{23} + \cdots + 1$. However all coefficients of the Alexander polynomial of an L -space knot are ± 1 .

Krcatovich [9] proved that all L -space knots are prime, so such connected sums are definitely not L -space knots. A proof that such connected sums cannot be *concordant* to L -space knots appears to be inaccessible using Heegaard Floer theory alone. The proof of the main theorem depends on a detailed analysis of the signature functions and Alexander polynomials of connected sums of torus knots. This dependence on classical invariants seems to be necessary, as the following example shows. We will see that the connected sum $T(2, 3) \# T(2, 3)$ is not concordant to an L -space knot. However, the torus knot $T(2, 5)$ is an L -space knot. The Heegaard Floer complex $\text{CFK}^\infty(T(2, 5))$ is formed from $\text{CFK}^\infty(T(2, 3) \# T(2, 3))$ by adding an acyclic summand, and thus known concordance invariants that arise from $\text{CFK}^\infty(K)$ cannot alone prove that K is not concordant to an L -space knot. (See the survey [5] for a discussion of the role of acyclic summands in Heegaard Floer knot theory.)

References. Basic facts about the Alexander polynomials of torus knots are covered in textbooks on knot theory, such as [3, 17]. For facts about L -space knots and the necessary Heegaard Floer theory, see [14, 15, 16]. Basic results concerning the Levine–Tristram signature function [10, 18], a step-function on the unit interval, are contained in the original sources. Its behavior under cabling is described in [11].

There is a jump function, $J_K(t)$, associated to the signature function of a knot; this is defined in Section 2.2. One fact about $J_K(t)$ that is used is that for each t , $|J_K(t)|$ is bounded above by the order of $e^{2\pi it}$ as a root of the Alexander polynomial $\Delta_K(t)$. Also, the jump equals the order of that root modulo two. This follows most easily from Milnor’s description [13] of what are now called *Milnor signatures*, which Matumoto [12] proved equal the jumps in the signature function; for recent presentations, see [6, 8].

Acknowledgments. Thanks are due to John Baldwin, Matt Hedden, Jen Hom, David Krcatovich, Adam Levine, and Ian Zemke. Valuable comments from the referee are also appreciated.

2. Torus knot Alexander polynomials and signature functions

2.1. Alexander polynomials of torus knots. The Alexander polynomial of the positive torus knot $T(p, q)$ is given by

$$(1) \quad \Delta_{p,q}(t) = \frac{(t^{pq} - 1)(t - 1)}{(t^p - 1)(t^q - 1)}.$$

Roots of this are pq -roots of unity that are not p -roots of unity or q -roots of unity. Letting $\phi_n(t)$ denote the n th cyclotomic polynomial, we have:

Lemma 2.1. *With notation as above,*

$$\Delta_{p,q}(t) = \prod \phi_{\alpha_j \beta_j}(t),$$

where the product is over the set of all pairs (α_j, β_j) for which α_j is a factor of p , β_j is a factor of q , and both are greater than 1.

Lemma 2.2. *Consider a set of n torus knots, $\{T(p_i, q_i)\}$, and let $d_i = (p_i - 1)(q_i - 1)$ be the degree of the $\Delta_{T(p_i, q_i)}(t)$. For $K = \#_i T(p_i, q_i)$, the two highest degree terms of the Alexander polynomial are $t^{\sum d_i} - nt^{(\sum d_i) - 1}$.*

Proof. Consider the numerator of the product of the Alexander polynomials when is written in the quotient form described by Equation (1). The leading term arises as the product of terms of degree $\sum(p_i q_i + 1)$. (This results from the product of the $t^{p_i q_i}$ terms times the product of the t terms in the n factors $(t - 1)$.) The next term, of degree one less, is obtained from a similar product, except one of the t terms from a $(t - 1)$ factor is not included, being replaced with -1 in the product. There are n such possible terms to drop.

The leading term of the denominator is degree $s = \sum p_i + \sum q_i$. The next highest degree term is of degree at most $s - 2$ (which would occur if one of the p_i or q_i were equal to two). The result stated in the lemma is now easily seen, for instance by considering the long division algorithm. \square

2.2. Signature functions of torus knots. For a knot $K \subset S^3$, let $\sigma_K(t)$ denote the signature of the hermitian form $(1 - \omega)V_K + (1 - \bar{\omega})V_K^T$, where V_K is a Seifert matrix and $\omega = e^{2\pi i t}$. The associated jump function is given by

$$J_K(t) = \frac{1}{2} \left(\lim_{s \rightarrow t^+} \sigma_K(s) - \lim_{s \rightarrow t^-} \sigma_K(s) \right).$$

This function is a concordance invariant of K . Figure 1 illustrates the graph of the signature function for the knot $T(3, 7)$ on $[0, \frac{1}{2}]$, which has jumps at $\{\frac{1}{21}, \frac{2}{21}, \frac{4}{21}, \frac{5}{21}, \frac{8}{21}, \frac{10}{21}\}$, each of value ± 1 . Notice that the jump at $\frac{1}{21}$ is negative. (The factor of $1/2$ in the definition of $J_K(t)$ is included to simplify notation. As defined, $J_K(t) = \sigma_K(t + \epsilon) - \sigma_K(t)$ for all sufficiently small $\epsilon > 0$.)

Lemma 2.3. *For any positive torus knot $T(p, q)$ and for any positive integer r , $J_K(1/r) \leq 0$.*

Proof. We use a formula of Litherland [11] to study the signature function. In this formula, we denote the integer lattice by Λ . For $0 \leq x \leq 1$, let

$$S_-(x) = \left\{ (i, j) \in \Lambda \mid 0 < i < p, 0 < j < q, \text{ and } \frac{i}{p} + \frac{j}{q} = x \right\}$$




FIGURE 1.

and

$$S_+(x) = \left\{ (i, j) \in \Lambda \mid 0 < i < p, 0 < j < q, \text{ and } \frac{i}{p} + \frac{j}{q} = 1 + x \right\}.$$

According to [11], $J_{T(p,q)}(x)$ is given by the difference of counts:

$$J_{T(p,q)}(x) = \#S_-(x) - \#S_+(x).$$

The lemma is a consequence of the observation that for any positive integer r , $\#S_-(1/r) = 0$. To see this, we consider the equation

$$\frac{i}{p} + \frac{j}{q} = \frac{1}{r}.$$

Multiplying by pq gives

$$iq + jp = \frac{pq}{r}.$$

Jumps in the signature function can occur only at roots of the Alexander polynomial. Applying Lemma 2.1, we need to consider the case of $r = \alpha_1\beta_1$, where $p = \alpha_1\alpha_2$, $\alpha_1 > 1$, $q = \beta_1\beta_2$, and $\beta_1 > 1$. Thus, our equation becomes

$$iq + jp = \alpha_2\beta_2.$$

Since α_2 divides p and $\alpha_2\beta_2$ and is relatively prime to q , it must also divide i . We write $i = i'\alpha_2$ and then divide by α_2 to find

$$i'q + j\alpha_1 = \beta_2.$$

Similarly, β_2 divides q (and itself), so it divides $j\alpha_1$. However, β_2 and α_1 are relatively prime, so $j = j'\beta_2$. Dividing yields

$$i'\beta_1 + j'\alpha_1 = 1.$$

Clearly, since both summands are at least one, the sum is at least two. Thus, there is no solution to the equation, and $S_-(1/r) = 0$, as desired. \square

3. Concordances to L -space knots

We now prove our main theorem.

Theorem 3.1. *Let $\{(p_i, q_i)\}_{i=1, \dots, n}$ be a set of pairs of relatively prime positive integers with $2 \leq p_i < q_i$ for all i and with $n > 1$. Then $\#_i T(p_i, q_i)$ is not concordant to an L -space knot.*

Proof. Let $K = \#_i T(p_i, q_i)$. Then $\Delta_K(t) = \prod_j \phi_{r_j}(t)^{m_j}$ for some set of distinct r_j and exponents $m_j > 0$. Notice that m_j is precisely the number of pairs (p_i, q_i) for which r_j can be written as a product of factor of p_i and a factor of q_i , both of which are greater than 1.

For those $T(p_i, q_i)$ for which $\phi_{r_j}(t)$ is a factor of $\Delta_{p_i, q_i}(t)$, that factor has exponent one, and hence the jump at $1/r_j$ is either ± 1 . By Lemma 2.3, the jump is -1 . It now follows that the jump in the signature function of K at $t = \frac{1}{r_j}$ equals $-m_j$.

Suppose that J is concordant to K . Then the jump in the signature function of J at $t = \frac{1}{r_j}$ also equals $-m_j$. Thus, $\phi_{r_j}(t)^{m_j}$ is a factor of $\Delta_J(t)$. It follows that $\text{degree}(\Delta_J(t)) \geq \text{degree}(\Delta_K(t))$.

For a connected sum of positive torus knots, $2\tau(K) = \text{degree}(\Delta_K(t))$. Also, for any L -space knot, $2\tau(J) = \text{degree}(\Delta_J(t))$. Thus, we have the inequalities

$$2\tau(K) = \text{degree}(\Delta_K(t)) \leq \text{degree}(\Delta_J(t)) = 2\tau(J).$$

But $\tau(J) = \tau(K)$, so we conclude

$$\Delta_K(t) = \Delta_J(t).$$

By Lemma 2.2, this polynomial does not have all nonzero coefficients equal to ± 1 , and thus it cannot be the Alexander polynomial of an L -space knot. \square

4. Generalizations

There are cases in which the main theorem extends to connected sums of torus knots, not all of which are positive. Here we discuss a few basic examples. Recently, Samantha Allen [1] proved that the subgroup of the concordance group generated by a pair of torus knots contains no nontrivial L -space knots other than the two torus knots themselves.

The simplest example is $K = T(2, 5) \# -T(2, 3)$. It has $\tau(K) = 1$ and Alexander polynomial $\phi_6(t)\phi_{10}(t)$. Since the signature function jumps at all the 6 and 10 roots of unity, any J concordant to K would have its Alexander polynomial divisible by $\Delta_K(t)$, and thus would have degree greater than 2. In particular since this degree exceeds twice the tau invariant, J could not be an L -space knot.

A second example is $K = -T(3, 4) \# -T(4, 5) \# T(5, 6)$. For this knot, $\tau(K) = 2$, and so if it were concordant to an L -space knot J , the degree of

$\Delta_J(t)$ would be four. This is impossible, since a calculation shows that the signature function of K has 32 singular points on the unit circle.

It is a simple matter to build a large collection of examples.

The first example for which results above do not apply is the connected sum of torus knots $K = T(2, 9) \# -T(2, 3)$. This knot has $\tau(K) = 3$ and Alexander polynomial

$$\Delta_K(t) = \phi_{18}(t)\phi_6(t)^2 = (1 - t^3 + t^6)(1 - t + t^2)^2.$$

The signature function jumps by one at each of the roots of ϕ_{18} , and only at those roots. Thus, the results proved above cannot rule out the possibility that K is concordant to an L -space knot J with $\Delta_J(t) = \phi_{18}(t)$.

In fact, deeper results from Heegaard Floer knot theory can be applied to the knot $T(2, 9) \# -T(2, 3)$. A theorem of Hedden and Watson [7, Corollary 9] states that for an L -space knot of genus g , the leading terms of the Alexander polynomial are $t^{2g} - t^{2g-1}$. (This also follows from a recent result of Baldwin and Vela-Vick [2] which states that if K is fibered of genus g , then $\widehat{\text{HFK}}(K, g - 1) \neq 0$.) However, $\phi_{18}(t) = t^6 - t^3 + 1$, which is not consistent with this constraint. Arguments along these lines yield large families of examples, but are not sufficient to give a general independence result. See [1] for much more general result of this type.

Another interesting area for extending the main theorem is that of algebraic knots. These can be described as iterated cables of torus knots of the form $T(p_1, q_1)_{(p_2, q_2), (p_3, q_3), \dots}$, for which all p_i and q_i are nonnegative and $q_{i+1} > p_i q_i p_{i+1}$ for all i . (A basic reference for algebraic knots is [4]. The algebraic properties of such knots are developed in [11].) The methods of this paper apply to show that many connected sums of algebraic knots are not concordant to L -space knots, but a general result has not been attained. Thus, we end with a question.

Question: Can a nontrivial connected sum of algebraic knots be concordant to an L -space knot?

References

- [1] ALLEN, SAMANTHA. Concordances from differences of torus knots to L -space knots. arXiv:1710.10664.
- [2] BALDWIN, JOHN; VELA-VICK, DAVID. A note on the knot Floer homology of fibered knots. arXiv:1801.06563.
- [3] BURDE, GERHARD; ZIESCHANG, HEINER. Knots. Second edition. De Gruyter Studies in Mathematics, 5. *Walter de Gruyter & Co., Berlin*, 2003. xii+559 pp. MR1959408 (2003m:57005), Zbl 1009.57003.
- [4] EISENBUD, DAVID; NEUMANN, WALTER. Three-dimensional link theory and invariants of plane curve singularities. *Annals of Mathematics Studies*, 110. *Princeton University Press*, 1985. vii+173 pp. MR0817982 (87g:57007), Zbl 0628.57002.
- [5] HOM, JENNIFER. A survey on Heegaard Floer homology and concordance. *J. Knot Theory Ramifications* **26** (2017) 1740015, 24pp. MR3604497, Zbl 1360.57002, arXiv:1512.00383, doi: 10.1142/S0218216517400156.

- [6] GILMER, PATRICK M.; LIVINGSTON, CHARLES. Signature jumps and Alexander polynomials for links. *Proc. Amer. Math. Soc.* **144** (2016), 5407–5417. MR3556282, Zbl 1361.57010, arXiv:1508.04394, doi:10.1090/proc/13129.
- [7] HEDDEN, MATTHEW; WATSON, LIAM. On the geography and botany of knot Floer homology. To appear, *Math. Selecta*. arXiv:1404.6913.
- [8] KEARNEY, M. KATE. The stable concordance genus. *New York J. Math.* **20** (2014), 973–987. MR3291607, Zbl 1331.57009, arXiv:1310.2513, <http://nyjm.albany.edu/j/2014/20-47.html>.
- [9] KRCATOVICH, DAVID. The reduced knot Floer complex. *Topology Appl.* **194** (2015) 171–201. MR3404612, Zbl 1326.57020, arXiv:1310.7624, doi:10.1016/j.topol.2015.08.008.
- [10] LEVINE, JEROME P. Invariants of knot cobordism. *Invent. Math.* **8** (1969) 98–110. MR0253348 (40 #6563), Zbl 0179.52401, doi:10.1007/BF01404613.
- [11] LITHERLAND, RICHARD A. Signatures of iterated torus knots. *Topology of Low-Dimensional Manifolds* (Proc. Second Sussex Conf., Chelwood Gate, 1977), 71–84. Lecture Notes in Math., 722. Springer, Berlin, 1979. MR0547456 (80k:57012), Zbl 0412.57002.
- [12] MATUMOTO, TAKAO. On the signature invariants of a non-singular complex sesquilinear form. *J. Math. Soc. Japan* **29** (1977) 67–71. MR0437456 (55 #10386), Zbl 0376.15013, doi:10.2969/jmsj/02910067.
- [13] MILNOR, JOHN. Infinite cyclic coverings. 1968 *Conference on the Topology of Manifolds* (Michigan State Univ., E. Lansing, Mich., 1967), 115–133. Prindle, Weber & Schmidt, Boston, Mass. MR0242163 (39 #3497), Zbl 0179.52302.
- [14] OZSVÁTH, PETER; SZABÓ, ZOLTÁN. Knot Floer homology and the four-ball genus. *Geom. Topol.* **7** (2003) 615–639. MR2026543 (2004i:57036), Zbl 1037.57027, arXiv:math.GT/0301149, doi:10.2140/gt.2003.7.615.
- [15] OZSVÁTH, PETER; SZABÓ, ZOLTÁN. Holomorphic disks and knot invariants. *Adv. Math.* **186** (2004), 58–116. MR2065507 (2005e:57044), Zbl 1062.57019, arXiv:math/0209056, doi:10.1016/j.aim.2003.05.001.
- [16] OZSVÁTH, PETER; SZABÓ, ZOLTÁN. On knot floer homology and lens space surgeries. *Topology* **44** (2005), 1281–1300. MR2168576 (2006f:57034), Zbl 1077.57012, arXiv:math/0303017, doi:10.1016/j.top.2005.05.001.
- [17] ROLFSEN, DALE. Knots and links, Mathematics Lecture Series, 7. *Publish or Perish, Inc., Berkeley, Calif.*, 1976. MR1277811 (95c:57018), Zbl 0854.57002.
- [18] TRISTRAM, A. Some cobordism invariants for links. *Proc. Camb. Phil. Soc.* **66** (1969), 251–264. MR0248854 (40 #2104), Zbl 0191.54703.
- [19] ZEMKE, IAN. Connected sums and involutive knot Floer homology. arXiv:1705.01117.

(Charles Livingston) DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405

livingst@indiana.edu

This paper is available via <http://nyjm.albany.edu/j/2018/24-12.html>.