

Sub-Hilbert relation for Fock–Sobolev type spaces

Setareh Eskandari, Ali Abkar, Per Åhag
and Antti Perälä

ABSTRACT. In this paper, two specific sub-Hilbert spaces are studied. They arise from the action of a Toeplitz operator on Fock–Sobolev type spaces, induced by a general Gaussian type weight. The argument is based on analysing the reproducing kernel of the corresponding sub-Hilbert space.

CONTENTS

1. Introduction	958
2. Fock-Sobolev type spaces	960
3. Gamma function and reproducing kernels	962
4. The main results	964
Acknowledgements	968
References	968

1. Introduction

This paper is concerned with sub-Hilbert functional spaces of analytic functions on planar domains. Suppose T is a bounded operator on a given Hilbert space H . We denote by $\mathcal{M}(T)$ the range of T , which is equipped with the following inner product:

$$\langle Tx, Ty \rangle_{\mathcal{M}(T)} = \langle x, y \rangle_H \quad x, y \in H \ominus \ker T.$$

Then $\mathcal{M}(T)$ is a Hilbert space. If, in addition, T is a contraction operator, the Hilbert space

$$\mathcal{M}((I - TT^*)^{1/2})$$

is called the complemented space to $\mathcal{M}(T)$ is denoted by $\mathcal{H}(T)$ and is called a sub-Hilbert space.

Received April 5, 2022.

2020 *Mathematics Subject Classification*. Primary 47B35; Secondary 30H05, 46E20.

Key words and phrases. Sub-Fock Hilbert space, Fock–Sobolev spaces, Toeplitz operator, reproducing kernel.

The pioneering work on sub-Hilbert spaces was done by L. de Branges, J. Rovnyak and D. Sarason [8, 9, 10, 18]. For further reading on the spaces introduced by de Branges and Rovnyak, their equivalent formulations, and their applications in function theory and operator theory, see [3]. Sarason’s monograph [18] contains extensive investigation of sub-Hilbert spaces arising from Toeplitz operators T_f acting on the Hardy space on the unit circle; in this context, it is customary to agree on the notation $\mathcal{M}(T_f) = \mathcal{M}(f)$ and $\mathcal{H}(T_f) = \mathcal{H}(f)$.

Later, continuing Sarason’s work, Kehe Zhu introduced sub-Bergman Hilbert spaces on the unit disk [21, 22]. To provide a brief account on this issue, we recall that the standard weighted Bergman space A_α^2 , for $\alpha > -1$, consists of all analytic functions on the unit disk for which the integral

$$\int_{\mathbb{D}} |f(z)|^2 (1 - |z|^2)^\alpha dx dy$$

is finite. The norm of a function in the weighted Bergman space is given by

$$\|f\|^2 = \frac{\alpha + 1}{\pi} \int_{\mathbb{D}} |f(z)|^2 (1 - |z|^2)^\alpha dx dy.$$

We shall at times write

$$dA_\alpha(z) = \frac{\alpha + 1}{\pi} (1 - |z|^2)^\alpha dx dy,$$

for normalized weighted area measure in the unit disk. Note that A_α^2 is a reproducing kernel functional Hilbert space whose kernel is given by

$$K_z^\alpha(w) = \sum_{n=0}^\infty \frac{\Gamma(n + \alpha + 2)}{n! \Gamma(\alpha + 2)} (\bar{z}w)^n = \frac{1}{(1 - w\bar{z})^{\alpha+2}}, \quad (z, w) \in \mathbb{D} \times \mathbb{D}.$$

The Bergman projection

$$P_\alpha : L^2(\mathbb{D}, dA_\alpha) \rightarrow A_\alpha^2(\mathbb{D})$$

is defined by

$$P_\alpha f(z) = \int_{\mathbb{D}} f(w) \overline{K_z^\alpha(w)} dA_\alpha(w).$$

Now, let φ be an analytic function in the unit disk satisfying $\|\varphi\|_\infty \leq 1$. For $\alpha \geq 0$, we consider the Toeplitz operator

$$T_\varphi^\alpha(f) = P_\alpha(\varphi f), \quad f \in A_\alpha^2.$$

For $\alpha = 0$, the unweighted Bergman space, Kehe Zhu [21, 22] studied the sub-Bergman Hilbert spaces $\mathcal{H}_\alpha(\varphi) := \mathcal{H}(T_\varphi^\alpha)$ and $\mathcal{H}_\alpha(\overline{\varphi}) := \mathcal{H}(T_\varphi^\alpha)$. He proved that these sub-Bergman Hilbert spaces coincide as sets, moreover, both spaces contain the Banach space of all bounded analytic functions on the unit disk. Zhu further showed that for the symbol z^m , and more generally, for a finite Blaschke product B , we have

$$\mathcal{H}(B) = \mathcal{H}(\overline{B}) = H^2,$$

where H^2 denotes the Hardy space of analytic functions on the unit disk.

Later, in 2010, Abkar and Jafarzade [1] extended Zhu's results to the standard weighted Bergman spaces A_α^2 where $\alpha \geq 0$. They proved that $H^\infty \subset \mathcal{H}_\alpha(\varphi) = \mathcal{H}_\alpha(\overline{\varphi})$, and for a finite Blaschke product B ,

$$H^\infty \subset \mathcal{H}_\alpha(B) = \mathcal{H}_\alpha(\overline{B}) = A_{\alpha-1}^2.$$

In 2014, this line of investigation was adapted by Nowak and Rososzczuk in [15] where the authors extended the latter result for $-1 < \alpha < 0$. They proved that

$$\mathcal{H}_\alpha(B) = \mathcal{H}_\alpha(\overline{B}) = \mathcal{D}_{\alpha+1},$$

where the Dirichlet space \mathcal{D}_α consists of all analytic functions f in the unit disk such that $f' \in L^2(\mathbb{D}, dA_\alpha)$. See also [20], and [16] where in the latter the authors studied similar problems in the unit ball of n -dimensional complex space \mathbb{C}^n .

Inspired by the aforementioned works, we will study the concept of a sub-Hilbert space in the context of Fock-type spaces $F_{\alpha,\beta,s}^2$, where the indices α and β appear in the exponential part of the weight, and s can be thought of as the order of the fractional derivative; see the next section. However, on these spaces, multiplication by an entire non-constant function is never bounded, let alone contractive. We will therefore focus our attention to the symbols of the type $f(z) = (z/|z|)^m$. We prove

Theorem 1. *Let $\alpha, \beta > 0$, $s \in \mathbb{R}$ and $m \in \mathbb{N}$, and let $T_f^{\alpha,\beta,s}$ be the Toeplitz operator on $F_{\alpha,\beta,s}^2$ induced by the symbol $f(z) = (z/|z|)^m$. We then have*

$$\mathcal{H}(f) = \mathcal{H}(\overline{f}) = F_{\alpha,\beta,s+\beta/2}^2.$$

2. Fock-Sobolev type spaces

Let \mathbb{C} denote the complex plane, $H(\mathbb{C})$ the space of entire functions, and $dA(z)$ the Lebesgue area measure on \mathbb{C} ;

$$dA(z) = \frac{1}{\pi} dx dy, \quad z = x + iy.$$

For $\alpha, \beta > 0$ and $s \in \mathbb{R}$, we consider the weight

$$d\lambda_{\alpha,\beta,s}(z) = |z|^{2s} e^{-\alpha|z|^\beta} dA(z).$$

In the literature, it is common to normalize $d\lambda_{\alpha,\beta,s}$ into a probability measure. However, when $s \leq -1$, this weight is no longer integrable, and cannot be normalized in an obvious way. We refrain from normalizing the weight altogether because of this.

2.1. Case $s > -1$. We define the generalized Fock-Sobolev type space $F^2_{\alpha,\beta,s}$ as those elements in $H(\mathbb{C})$ that are square integrable over \mathbb{C} with respect to $d\lambda_{\alpha,\beta,s}$. That is,

$$F^2_{\alpha,\beta,s} = L^2_{\alpha,\beta,s} \cap H(\mathbb{C}).$$

It is easy to see that $F^2_{\alpha,\beta,s}$ is a closed subspace of $L^2_{\alpha,\beta,s} = L^2(\mathbb{C}, d\lambda_{\alpha,\beta,s})$, and a Hilbert space with the inner product.

$$\langle f, g \rangle_{\alpha,\beta,s} = \int_{\mathbb{C}} f(z)\overline{g(z)}d\lambda_{\alpha,\beta,s}(z).$$

2.2. Case $s \leq -1$. The spaces $F^2_{\alpha,\beta,s}$ also make sense for $s \leq -1$, but in that case, following the definition above would require the members $F^2_{\alpha,\beta,s}$ to have a deep enough zero at the origin. In [6] two ways to overcome this are presented. First, one could replace the term $|z|^{2s}$ in $d\lambda_{\alpha,\beta,s}$ by $(1 + |z|)^{2s}$. However, the other approach from [6] fits our calculations better. Given

$$f(z) = \sum_{k=0}^{\infty} f_k z^k,$$

let us denote by $p_N(f)$ the degree N Maclaurin polynomial of f ;

$$p_N(f)(z) = \sum_{k=0}^N f_k z^k.$$

Then, denote by $R_N(f) = f - p_N(f)$ the remainder, which in our case is going to determine the membership in $F^2_{\alpha,\beta,s}$.

By using the ceiling function, we define $N = -[s] - 1$ and introduce the inner product

$$\langle f, g \rangle_{\alpha,\beta,s} = \int_{\mathbb{C}} R_N(f)(z)\overline{R_N(g)(z)}d\lambda_{\alpha,\beta,s}(z) + \sum_{k=0}^N f_k \overline{g_k}.$$

The space $F^2_{\alpha,\beta,s}$ consists of entire functions f with

$$\|f\|^2_{\alpha,\beta,s} := \langle f, f \rangle_{\alpha,\beta,s} < \infty,$$

and by the virtue of the above definition, always contains all polynomials. In practice, we will not need to worry about this definition, as we are only interested in $R_N(f)$ for large enough N .

2.3. Relation to other Fock spaces. For particular choice of parameters, the spaces $F^2_{\alpha,\beta,s}$ reduce to more well-known spaces. The choice $(\alpha, \beta, s) = (\alpha, 2, 0)$ gives rise to classical Fock spaces, where standard references include the book of Folland [11] and the more recent book of Zhu [23].

Adding the parameter s is known to be equivalent to the membership of (fractional) derivatives in the standard Fock space. This motivates the terminology

Fock-Sobolev space, which corresponds to the choice $(\alpha, \beta, s) = (\alpha, 2, s)$ studied in [7, 6, 5]. These references do not always contain α as a parameter, but passage to this more general case is easy for most purposes of this paper.

In [4], Bommier-Hato, Engliš and Youssfi studied the so-called Fock-type spaces. These correspond to changing the Gaussian in the weight: $(\alpha, \beta, s) = (1, \beta, 0)$. Here again, slightly more general parameters do not cause much of an obstacle.

Finally, there are several generalization of the Fock-spaces to the case where the weight is non-radial; we mention [12], [14] and [19], but there are many more. These spaces are often called generalized Fock spaces, but we refrain from studying them, because having a radial weight is essential for our approach.

3. Gamma function and reproducing kernels

3.1. Gamma function. The Euler Gamma function (or simply the Gamma function) is a well-known special function that generalizes the concept of a factorial to non-integer values. As we have already seen, it appears naturally in the context of exponential weights.

The Gamma function can be defined by a convergent improper integral:

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt, \quad \operatorname{Re}(z) > 0.$$

The Gamma function satisfies the crucial recurrence relation: $\Gamma(z+1) = z\Gamma(z)$, and the following standard estimate for fixed complex numbers a and b

$$\frac{\Gamma(z+a)}{\Gamma(z+b)} \asymp z^{a-b}, \quad z \rightarrow \infty.$$

In this paper, we will need a more refined variant of the latter. The following formula can be found in [13].

$$\begin{aligned} \frac{\Gamma(z+a)}{\Gamma(z+b)} &= z^{a-b} \left[1 + \frac{(a-b)(a+b-1)}{2z} + \frac{(a-b)(a-b-1)}{24z^2} \right. \\ &\quad \left. \times \{3(a+b-1)^2 - a + b - 1\} \right] [1 + O(z^{-3})], \quad z \rightarrow \infty. \quad (1) \end{aligned}$$

Lemma 2. *The Gamma function satisfies*

$$1 - \frac{(\Gamma(z + \frac{a+b}{2}))^2}{\Gamma(z+a)\Gamma(z+b)} = \frac{(a-b)^2}{4z} + O(z^{-2}), \quad z \rightarrow \infty.$$

Proof. By using the equation (1) we can obtain estimates for $\frac{\Gamma(z + \frac{a+b}{2})}{\Gamma(z+a)}$ and $\frac{\Gamma(z + \frac{a+b}{2})}{\Gamma(z+b)}$, so that we have

$$\frac{\Gamma(z + \frac{a+b}{2})}{\Gamma(z+a)} = z^{\frac{b-a}{2}} \left[1 + \frac{(b-a)(3a+b-2)}{8z} + O(z^{-2}) \right]$$

and

$$\frac{\Gamma(z + \frac{a+b}{2})}{\Gamma(z + b)} = z^{\frac{a-b}{2}} \left[1 + \frac{(a-b)(a+3b-2)}{8z} + O(z^{-2}) \right]$$

as $z \rightarrow \infty$. Multiplying these completes the proof. □

The equation (1) can also be used to partially refine the recurrence relation:

Lemma 3. *For any complex number δ the Gamma function satisfies*

$$\Gamma(z + \delta) \asymp z^\delta \Gamma(z), \quad z \rightarrow \infty.$$

Proof. The proof is easy and we omit the details. □

3.2. Reproducing kernels and projections. The approach of this paper is based on identifying the sub-Hilbert space by calculating its reproducing kernel. This is a well-known approach, see [1, 18, 21, 22]. The theory of reproducing kernels is a fascinating field in its own right, extending far beyond what is needed here. Some classical references include [2] and [17].

Since the weight $d\lambda_{\alpha,\beta,s}$ is radial, the Fock-type space $F^2_{\alpha,\beta,s}$ possesses a monomial Schauder basis. If $s \leq -1$ and $n \leq -[s] - 1$, we set $e_n^{\alpha,\beta,s}(z) = z^n$ and observe that $\|e_n^{\alpha,\beta,s}\|_{\alpha,\beta,s} = 1$. Otherwise, we compute in polar coordinates and using the change of variables $t = \alpha r^\beta$:

$$\begin{aligned} \|z^n\|_{\alpha,\beta,s}^2 &= \int_{\mathbb{C}} |z|^{2n} |z|^{2s} e^{-\alpha|z|^\beta} dA(z) \\ &= \frac{1}{\pi} \int_0^\infty \int_0^{2\pi} r^{2n+2s+1} e^{-\alpha r^\beta} d\theta dr \\ &= \frac{2}{\beta \alpha^{\frac{2n+2s+2}{\beta}}} \int_0^\infty t^{\frac{2n+2s+2}{\beta}-1} e^{-t} dt \\ &= \frac{2}{\beta \alpha^{\frac{2n+2s+2}{\beta}}} \Gamma\left(\frac{2n+2s+2}{\beta}\right). \end{aligned}$$

So, for $n > -[s] - 1$, we observe that then the functions

$$e_n^{\alpha,\beta,s} = \sqrt{\frac{\beta \alpha^{\frac{2n+2s+2}{\beta}}}{2\Gamma\left(\frac{2n+2s+2}{\beta}\right)}} z^n$$

are unit vectors, and $(e_n^{\alpha,\beta,s})_{n=0}^\infty$ forms the basis of $F^2_{\alpha,\beta,s}$.

Let $K_z^{\alpha,\beta,s}$ denote the reproducing kernel of $F^2_{\alpha,\beta,s}$ – that is, the unique function in $F^2_{\alpha,\beta,s}$ with the property

$$f(z) = \langle f, K_z^{\alpha,\beta,s} \rangle_{\alpha,\beta,s}, \quad f \in F^2_{\alpha,\beta,s}.$$

By a well-known identity, we obtain

$$K_z^{\alpha,\beta,s}(\xi) = \sum_{n=0}^{\infty} \frac{\beta \alpha^{\frac{2n+2s+2}{\beta}}}{2\Gamma\left(\frac{2n+2s+2}{\beta}\right)} (\xi \bar{z})^n,$$

when $s > -1$. When $s \leq -1$, we obtain

$$\begin{aligned} K_z^{\alpha,\beta,s}(\xi) &= \sum_{n=0}^{\infty} e_n^{\alpha,\beta,s}(\xi) \overline{e_n^{\alpha,\beta,s}(z)} \\ &= \frac{1 - (\xi \bar{z})^{-|s|}}{1 - \xi \bar{z}} + \sum_{n=-|s|}^{\infty} \frac{\beta \alpha^{\frac{2n+2s+2}{\beta}}}{2\Gamma\left(\frac{2n+2s+2}{\beta}\right)} (\xi \bar{z})^n. \end{aligned}$$

In either case, we are only interested the asymptotics of the general term in the sum as n is large; that is

$$\frac{\beta \alpha^{\frac{2n+2s+2}{\beta}}}{2\Gamma\left(\frac{2n+2s+2}{\beta}\right)} \asymp \frac{\alpha^{\frac{2n}{\beta}}}{\Gamma\left(\frac{2n+2s+2}{\beta}\right)}. \tag{2}$$

In general, these power series can be understood in terms of the generalized Mittag-Leffler functions; see [4]. Of course, it is well-known (there are many references, see for instance [23]) that

$$K_z^{\alpha,2,0}(\xi) = \alpha e^{\alpha \xi \bar{z}}.$$

Finally, we are now able to write the orthogonal projection (the Bergman projection) $P^{\alpha,\beta,s} : L^2_{\alpha,\beta,s} \rightarrow F^2_{\alpha,\beta,s}$ as

$$P^{\alpha,\beta,s} f(z) = \int_{\mathbb{C}} f(\xi) \overline{K_z^{\alpha,\beta,s}(\xi)} d\lambda_{\alpha,\beta,s}(\xi).$$

By the standard theory of orthogonal projections, $P^{\alpha,\beta,s}$ is bounded; in fact the norm of $P^{\alpha,\beta,s}$ is one.

4. The main results

4.1. Toeplitz operators. Before proving the main result, a short discussion on Toeplitz operators in in order. Given an essentially bounded function f on the complex plane, let M_f denote the multiplication induced by f . It is clearly bounded from $F^2_{\alpha,\beta,s} \rightarrow L^2_{\alpha,\beta,s}$. The Toeplitz operator

$$T_f^{\alpha,\beta,s} : F^2_{\alpha,\beta,s} \rightarrow F^2_{\alpha,\beta,s}$$

is then defined as

$$T_f^{\alpha,\beta,s} = P^{\alpha,\beta,s} M_f.$$

Observe that $T_f^{\alpha,\beta,s}$ is contractive, whenever $\|f\|_\infty \leq 1$. Since orthogonal projections are self-adjoint, it is easy to see that the adjoint of $T_f^{\alpha,\beta,s}$ equals $T_{\bar{f}}^{\alpha,\beta,s}$. In particular, if $\|f\|_\infty \leq 1$, the operators

$$I - T_f^{\alpha,\beta,s} T_{\bar{f}}^{\alpha,\beta,s} \quad \text{and} \quad I - T_{\bar{f}}^{\alpha,\beta,s} T_f^{\alpha,\beta,s}$$

are positive.

In [1, 18, 21, 22] the authors study function spaces on the unit disk, and problem of sub-Hilbert spaces induced by a Toeplitz operator (given by the orthogonal projection of the respective space). Special focus is given to symbols $f(z) = z^m$ and f being a finite Blaschke product. Neither option seems to work directly for our setting. Instead we take:

$$f(z) = \left(\frac{z}{|z|}\right)^m \quad \text{and} \quad \bar{f}(z) = \left(\frac{\bar{z}}{|z|}\right)^m,$$

where the contractivity requirement is automatically satisfied.

4.2. Proof of the main result. We will make use of the following result, which can be found in [18] (it is proven for the unit disk, but the exact same argument works for any reproducing kernel Hilbert space).

Lemma 4. *Let H be a reproducing kernel Hilbert space over a domain Ω , K_z its reproducing kernel and $T : H \rightarrow H$ a contraction. Then the reproducing kernel of $\mathcal{H}(T)$ is given by $(I - TT^*)K_z$.*

Note that every $F_{\alpha,\beta,s}^2$ is isometrically isomorphic to a weighted ℓ^2 space, with the weight coming from the moments of the weight $d\lambda_{\alpha,\beta,s}(z)$. On the other hand, also the reproducing kernel is related to these moments. Therefore, in order to determine $\mathcal{H}(T)$ and $\mathcal{H}(T^*)$, it suffices to study the asymptotic of the power series expansion of the reproducing kernel.

We are now in position to prove the main theorem. Let $\alpha, \beta > 0$ and $s \in \mathbb{R}$, and let $T_f^{\alpha,\beta,s}$ be the Toeplitz operator on $F_{\alpha,\beta,s}^2$ induced by the symbol $f(z) = (z/|z|)^m$. We then have

$$\mathcal{H}(f) = \mathcal{H}(\bar{f}) = F_{\alpha,\beta,s+\beta/2}^2.$$

We now prove this.

Proof. Suppose m is a natural number. We will calculate the reproducing kernels of sub-Fock-Sobolev Hilbert spaces. The formula

$$(I - T_{(\frac{\bar{z}}{|z|})^m} T_{(\frac{z}{|z|})^m}) K_z^{\alpha,\beta,s}$$

gives the reproducing kernels of these spaces. So, we consider the Toeplitz operator induced by $(\frac{z}{|z|})^m$ in Fock-Sobolev spaces $F_{\alpha,\beta,s}^2$. By using the definition

of Toeplitz operator and the formula (2), we have

$$\begin{aligned}
\frac{2}{\beta} \alpha^{\frac{-2s-2}{\beta}} T_{\left(\frac{z}{|z|}\right)^m} z^n &= \int_{\mathbb{C}} \left(\frac{\xi}{|\xi|}\right)^m \xi^n \sum_{k \geq 0} \frac{1}{\Gamma\left(\frac{2k+2s+2}{\beta}\right)} (\alpha^{2/\beta} z \bar{\xi})^k |\xi|^{2s} e^{-\alpha|\xi|^\beta} dA(\xi) \\
&= \frac{1}{\Gamma\left(\frac{2}{\beta}(n+m+s+1)\right)} \alpha^{\frac{2}{\beta}(m+n)} z^{m+n} \int_{\mathbb{C}} |\xi|^{2n+2s+m} e^{-\alpha|\xi|^\beta} dA(\xi) \\
&= \frac{\alpha^{\frac{2}{\beta}(m+n)}}{\Gamma\left(\frac{2}{\beta}(n+m+s+1)\right)} z^{m+n} 2 \int_0^\infty r^{2n+2s+m+1} e^{-\alpha r^\beta} dr \\
&= \frac{\alpha^{\frac{2}{\beta}(m+n)}}{\Gamma\left(\frac{2}{\beta}(n+m+s+1)\right)} z^{m+n} \frac{2}{\alpha\beta} \int_0^\infty \left(\frac{t}{\alpha}\right)^{\frac{1}{\beta}(2n+m+2s+2)-1} e^{-t} dt \\
&= \frac{\alpha^{\frac{1}{\beta}(m-2s-2)}}{\Gamma\left(\frac{2}{\beta}(n+m+s+1)\right)} z^{m+n} \frac{2}{\beta} \Gamma\left(\frac{2}{\beta}(n+s+1) + \frac{m}{\beta}\right) \\
&= \frac{\Gamma\left(\frac{2}{\beta}(n+s+1) + \frac{m}{\beta}\right)}{\Gamma\left(\frac{2}{\beta}(n+m+s+1)\right)} \frac{2}{\beta} z^{m+n} \alpha^{\frac{1}{\beta}(m-2s-2)}.
\end{aligned}$$

By a similar calculation for $T_{\left(\frac{z}{|z|}\right)^m} z^n$, using (2) we have

$$\begin{aligned}
\frac{2}{\beta} \alpha^{\frac{-2s-2}{\beta}} T_{\left(\frac{z}{|z|}\right)^m} z^n &= \int_{\mathbb{C}} \left(\frac{\bar{\xi}}{|\xi|}\right)^m \xi^n \sum_{k \geq 0} \frac{1}{\Gamma\left(\frac{2}{\beta}(k+s+1)\right)} (\alpha^{2/\beta} z \bar{\xi})^k |\xi|^{2s} e^{-\alpha|\xi|^\beta} dA(\xi) \\
&= \frac{\alpha^{\frac{2}{\beta}(n-m)}}{\Gamma\left(\frac{2}{\beta}(n-m+s+1)\right)} z^{n-m} \int_{\mathbb{C}} |\xi|^{2n+2s-m} e^{-\alpha|\xi|^\beta} dA(\xi) \\
&= \frac{\alpha^{\frac{2}{\beta}(n-m)}}{\Gamma\left(\frac{2}{\beta}(n-m+s+1)\right)} z^{n-m} 2 \int_0^\infty r^{2n+2s-m+1} e^{-\alpha r^\beta} dr \\
&= \frac{\alpha^{\frac{2}{\beta}(n-m)}}{\Gamma\left(\frac{2}{\beta}(n-m+s+1)\right)} z^{n-m} \frac{2}{\alpha\beta} \int_0^\infty \left(\frac{t}{\alpha}\right)^{\frac{1}{\beta}(2n-m+2s+2)-1} e^{-t} dt \\
&= \frac{\alpha^{\frac{1}{\beta}(-m-2s-2)}}{\Gamma\left(\frac{2}{\beta}(n-m+s+1)\right)} z^{n-m} \frac{2}{\beta} \Gamma\left(\frac{2}{\beta}(n+s+1) - \frac{m}{\beta}\right)
\end{aligned}$$

$$= \frac{\Gamma(\frac{2}{\beta}(n+s+1) - \frac{m}{\beta})}{\Gamma(\frac{2}{\beta}(n-m+s+1))} \frac{2}{\beta} z^{n-m} \alpha^{\frac{1}{\beta}(-m-2s-2)}.$$

It follows that

$$\left(I - T_{\left(\frac{z}{|z|}\right)^m} T_{\left(\frac{\bar{z}}{|z|}\right)^m} \right) z^n = \left(1 - \frac{\Gamma(\frac{2}{\beta}(n+s+1) + \frac{m}{\beta}) \Gamma(\frac{2}{\beta}(n+s+1) + \frac{m}{\beta})}{\Gamma(\frac{2}{\beta}(s+m+n+1)) \Gamma(\frac{2}{\beta}(s+n+1))} \right) z^n.$$

From Lemma (2), we conclude that

$$1 - \frac{\Gamma\left(\frac{2}{\beta}(n+s+1) + \frac{m}{\beta}\right) \Gamma\left(\frac{2}{\beta}(n+s+1) + \frac{m}{\beta}\right)}{\Gamma\left(\frac{2}{\beta}(s+m+n+1)\right) \Gamma\left(\frac{2}{\beta}(s+n+1)\right)} \asymp \frac{1}{n},$$

therefore

$$\left(I - T_{\left(\frac{\bar{z}}{|z|}\right)^m} T_{\left(\frac{z}{|z|}\right)^m} \right) K_z^{\alpha, \beta, s}(\xi) = \sum_{n=0}^{\infty} \frac{1}{n} \frac{1}{\Gamma\left(\frac{2n+2s+2}{\beta}\right)} (\alpha^{2/\beta} \xi \bar{z})^n. \tag{3}$$

For large enough n , we have

$$\begin{aligned} n \Gamma\left(\frac{2n+2s+2}{\beta}\right) &\asymp \left(\frac{2n+2s+2}{\beta}\right) \Gamma\left(\frac{2n+2s+2}{\beta}\right) \\ &= \Gamma\left(\frac{2n+2(s+\frac{\beta}{2})+2}{\beta}\right), \end{aligned} \tag{4}$$

Substituting (4) into (3), we get the reproducing kernel of Fock-Sobolev space of order $(s + \frac{\beta}{2})$, which completes the proof. \square

As a consequence of the main theorem, we obtain the following corollary for the Fock-Sobolev space $F_{\alpha, 2, s}^2$.

Corollary 5. *Let $f(z) = (z/|z|)^m$, and let us consider Toeplitz operators acting on $F_{\alpha, 2, s}^2$. Then*

$$\mathcal{H}(f) = \mathcal{H}(\bar{f}) = F_{\alpha, 2, s+1}^2.$$

Note that this is in line with the well-known Bergman space results of Zhu [21, 22] and Abkar-Jafarzadeh [1].

Acknowledgements

This paper was written when the first named author was visiting the Department of Mathematics and Mathematical Statistics of Umeå University during July 2021 to April 2022. She wishes to thank the chairman Prof. Åke Brännström and the whole department for all the warm hospitality.

References

- [1] ABKAR, ALI; JAFARZADEH, BAGHER. Weighted sub-Bergman Hilbert spaces in the unit disk. *Czechoslovak Math. J.* **60(135)** (2010), no. 2, 435–443. [MR2657960](#), [Zbl 1220.47033](#), doi: [10.1007/s10587-010-0026-2](#). [960](#), [963](#), [965](#), [967](#)
- [2] ARONSZAJN, NACHMAN. Theory of reproducing kernels. *Trans. Am. Math. Soc.* **68** (1950), 337–404. [MR0051437](#), [Zbl 0037.20701](#), doi: [10.1090/S0002-9947-1950-0051437-7](#). [963](#)
- [3] BALL, JOSEPH A.; BOLOTNIKOV, VLADIMIR. de Branges–Rovnyak spaces: basics and theory. in: *Operator Theory*, Alpay D. (Ed.), 631–679. *Springer, Basel*, 2015. [Zbl 1347.46018](#), doi: [10.1007/978-3-0348-0667-1_6](#). [959](#)
- [4] BOMMIER-HATO, HÉLÈNE; ENGLIŠ, MIROSLAV; YOUSSEFI, EL-HASSAN. Bergman-type projections in generalized Fock spaces. *J. Math. Anal. Appl.* **389** (2012), no. 2, 1086–1104. [MR2879282](#), [Zbl 1252.47023](#), [arXiv:1105.5600](#), doi: [10.1016/j.jmaa.2011.12.045](#). [962](#), [964](#)
- [5] CHO, HONG RAE; CHOE, BOO RIM; KOO, HYUNG WOO. Fock–Sobolev spaces of fractional order. *Potential Anal.* **43** (2015), no. 2, 199–240. [MR3374109](#), [Zbl 1327.32009](#), doi: [10.1007/s11118-015-9468-3](#). [962](#)
- [6] CHO, HONG RAE; ISRALOWITZ, JOSHUA; JOO, JAE CHEON. Toeplitz operators on Fock–Sobolev type spaces. *Integr. Equ. Opr. Theory* **82**, (2015), no. 1, 1–32. [MR3335506](#), [Zbl 1326.47029](#), doi: [10.1007/s00020-015-2223-8](#). [961](#), [962](#)
- [7] CHO, HONG RAE; ZHU, KEHE. Fock–Sobolev spaces and their Carleson measures. *J. Funct. Anal.* **263** (2012), no. 8, 2483–2506. [MR2964691](#), [Zbl 1264.46017](#), doi: [10.1016/j.jfa.2012.08.003](#). [962](#)
- [8] DE BRANGES, LOUIS; ROVNYAK, JAMES. Square summable power series. *Holt, Rinehart and Winston, New York-Toronto, Ont.-London*, 1966. viii+104 pp. [MR0215065](#), [Zbl 0153.39602](#). [959](#)
- [9] DE BRANGES, LOUIS; ROVNYAK, JAMES. Appendix on square summable power series. in: *Perturbation theory and its applications in quantum mechanics*, (C. Wilcox, ed.) 347–392. *John Wiley and Sons, New York*, 1966. [959](#)
- [10] DE BRANGES, LOUIS; ROVNYAK, JAMES. Canonical models in quantum scattering theory. in: *Perturbation theory and its applications in quantum mechanics* (C. Wilcox, ed.), 295–392. *John Wiley and Sons, New York*, 1966. [959](#)
- [11] FOLLAND, GERALD B. Harmonic analysis in phase space. *Annals of Mathematics Studies*, 122. *Princeton University Press, Princeton, NJ*, 1989. x+277 pp. ISBN: 0-691-08527-7; 0-691-08528-5. [MR0983366](#), [Zbl 0682.43001](#), doi: [10.1515/9781400882427](#). [961](#)
- [12] HU, ZHANGJIAN; VIRTANEN, JANI A. Schatten class Hankel operators on the Segal–Bargmann space and the Berger–Coburn phenomenon. *Trans. Amer. Math. Soc.* **375** (2022), no. 5, 3733–3753. [MR4402674](#), [Zbl 7502511](#), doi: [10.1090/tran/8638](#). [962](#)
- [13] LUKE, YUDELL L. The special functions and their approximations. I. *Mathematics in Science and Engineering*, 53. *Academic Press, New York-London*, 1969. xx+349 pp. [MR0241700](#), [Zbl 0193.01701](#). [962](#)
- [14] MARCO, NICOLAS; MASSANEDA, XAVIER; ORTEGA-CERDÁ, JOAQUIM. Interpolating and sampling sequences for entire functions. *Geom. Funct. Anal.* **13** (2003), no. 4, 862–914. [MR2006561](#), [Zbl 1097.30041](#), [arXiv:math/0205241](#), doi: [10.1007/s00039-003-0434-7](#). [962](#)

- [15] NOWAK, MARIA; ROSOSZCZUK, RENATA. Weighted sub-Bergman Hilbert spaces. *Ann. Univ. Mariae Curie-Skłodowska Sect. A* **68** (2014), no. 1, 49–57. [MR3252516](#), [Zbl 1290.30067](#), doi: [10.2478/umcsmath-2014-0006](#). [960](#)
- [16] ROSOSZCZUK, RENATA; SYMESAK, FRÉDÉRIC. Weighted sub-Bergman Hilbert spaces in the unit ball of \mathbb{C}^n . *Concr. Oper.* **7** (2020), no. 1, 124–132. [MR4151088](#), [Zbl 1453.32005](#), doi: [10.1515/conop-2020-0103](#). [960](#)
- [17] SAITOH, SABUROU. *Theory of Reproducing Kernels and its Applications*. Pitman Research Notes in Mathematics Series, 189. Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1988. x+157 pp. ISBN: 0-582-03564-3. [MR0983117](#), [Zbl 0652.30003](#). [963](#)
- [18] SARASON, DONALD. Sub-Hardy Hilbert spaces in the unit disk. University of Arkansas Lecture Notes in the Mathematical Sciences, 10. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1994. xvi+95 pp. ISBN: 0-471-04897-6. [MR1289670](#), [Zbl 1253.30002](#). [959](#), [963](#), [965](#)
- [19] SCHUSTER, ALEXANDER P; VAROLIN, DROR. Toeplitz operators and Carleson measures on generalized Bargmann–Fock spaces. *Integral Equations Operator Theory* **72** (2012), no. 3, 363–392. [MR2891634](#), [Zbl 1262.47047](#), doi: [10.1007/s00020-011-1939-3](#). [962](#)
- [20] SYMESAK, FRÉDÉRIC. Sub-Bergman spaces in the unit ball of \mathbb{C}^n . *Proc. Amer. Math. Soc.* **138** (2010), no. 12, 4405–4411. [MR2680064](#), [Zbl 1208.32006](#), doi: [10.1090/S0002-9939-2010-10437-9](#). [960](#)
- [21] ZHU, KEHE. Sub-Bergman Hilbert spaces on the unit disk. *Indiana Univ. Math. J.* **45** (1996), no. 1, 165–176. [MR1406688](#), [Zbl 0863.30051](#), doi: [10.1512/iumj.1996.45.1097](#). [959](#), [963](#), [965](#), [967](#)
- [22] ZHU, KEHE. Sub-Bergman Hilbert spaces in the unit disk, II. *J. Funct. Anal.* **202** (2003), no. 2, 327–341. [MR1990528](#), [Zbl 1039.47019](#), doi: [10.1016/S0022-1236\(02\)00086-1](#). [959](#), [963](#), [965](#), [967](#)
- [23] ZHU, KEHE. Analysis on Fock spaces. Graduate Texts in Mathematics, 263. Springer-Verlag, New York, 2012. x+344 pp. ISBN: 978-1-4419-8800-3. [MR2934601](#), [Zbl 1262.30003](#), doi: [10.1007/978-1-4419-8801-0](#). [961](#), [964](#)

(Setareh Eskandari) DEPARTMENT OF MATHEMATICS AND MATHEMATICAL STATISTICS, UMEÅ UNIVERSITY, 90187 UMEÅ, SWEDEN, AND DEPARTMENT OF PURE MATHEMATICS, FACULTY OF SCIENCE, IMAM KHOMEINI INTERNATIONAL UNIVERSITY, 34149 QAZVIN, IRAN
s.eskandari@edu.ikiu.ac.ir

(Ali Abkar) DEPARTMENT OF PURE MATHEMATICS, FACULTY OF SCIENCE, IMAM KHOMEINI INTERNATIONAL UNIVERSITY, 34149 QAZVIN, IRAN
abkar@sci.ikiu.ac.ir

(Per Åhag) DEPARTMENT OF MATHEMATICS AND MATHEMATICAL STATISTICS, UMEÅ UNIVERSITY, 90187 UMEÅ, SWEDEN
per.ahag@umu.se

(Antti Perälä) DEPARTMENT OF MATHEMATICS AND MATHEMATICAL STATISTICS, UMEÅ UNIVERSITY, 90187 UMEÅ, SWEDEN
antti.perala@umu.se

This paper is available via <http://nyjm.albany.edu/j/2022/28-39.html>.