

Twistor fibers in hypersurfaces of the flag threefold

**Amedeo Altavilla, Edoardo Ballico
and Maria Chiara Brambilla**

ABSTRACT. We study surfaces of bidegree $(1, d)$ contained in the flag threefold in relation to the twistor projection. In particular, we focus on the number and the arrangement of twistor fibers contained in such surfaces. First, we prove that there is no irreducible surface of bidegree $(1, d)$ containing $d+2$ twistor fibers in general position. On the other hand, given any collection of $(d+1)$ twistor fibers satisfying a mild natural constraint, we prove the existence of a surface of bidegree $(1, d)$ that contains them. We improve our results for $d=2$ or $d=3$, by removing all the generality hypotheses.

CONTENTS

1. Introduction	1117
2. Preliminaries and first results	1120
3. Surfaces of bidegree $(1, d)$	1129
4. Surfaces of bidegree $(1, 2)$ and $(1, 3)$	1135
References	1147

1. Introduction

The study of the twistor geometry of the flag threefold is motivated by the search for Riemannian 4-manifolds admitting several integrable complex structures compatible with the prescribed metric (see e.g. [8, 12, 13]). In this context, a recent trend is the investigation of specific cases in order to find explicit examples [1, 2, 5, 7, 9].

The flag threefold \mathbb{F} can be seen as the twistor space of the complex projective plane \mathbb{P}^2 endowed with all its standard structures [6, 11],

$$\pi : \mathbb{F} \rightarrow \mathbb{P}^2.$$

Received May 30, 2023.

2010 *Mathematics Subject Classification.* Primary: 32L25, 14M15; Secondary: 14D21, 32M10, 14J26.

Key words and phrases. Flag threefold, twistor projection, twistor fiber, surfaces, bidegree.

All the authors are partially supported by GNSAGA. The first named author is partially supported by the INdAM project ‘Teoria delle funzioni ipercomplesse e applicazioni’.

The threefold \mathbb{F} embeds in $\mathbb{P}^2 \times \mathbb{P}^2$, hence it is possible to define a natural notion of bidegree (d_1, d_2) for curves and surfaces in \mathbb{F} .

In [4] we have started a detailed analysis of the geometry of the algebraic curves and surfaces contained in the flag threefold \mathbb{F} , in relation to the twistor projection. In particular, twistor fibers are smooth irreducible curves of bidegree $(1,1)$. In [3] we gave a first bound on the maximum number of smooth irreducible curves of bidegree $(1, 1)$ contained in a smooth surface $S \subset \mathbb{F}$. Here, by focussing our attention on a particular family of surfaces, we obtain stronger and more significant results. In fact, we analyse the case of bidegree $(1, d)$ surfaces in \mathbb{F} and study the number and the arrangements of twistor fibers contained in them.

A surface S of bidegree $(1, d)$ can be realised as a $d : 1$ branched cover of \mathbb{P}^2 , or alternatively as a blow up of \mathbb{P}^2 in $n = 1 + d + d^2$ points. If $d = 1$ then S is a (toric) del Pezzo surface of degree 6, and this case was studied in [4]. In the present paper we give results for arbitrary d , as well as focussing on $d = 2$ and $d = 3$. In general, if S has bidegree (d_1, d_2) , the twistor projection π restricted to S is a branched cover of degree $d_1 + d_2$ of \mathbb{P}^2 , so if $(d_1, d_2) = (1, d)$, it is natural to compare our results with those obtained for degree $d + 1$ surfaces in \mathbb{P}^3 viewed as the twistor space of the 4-sphere [11]. In the latter case, surfaces of degree 2 and 3 have been studied in some detail. In particular, surfaces of degree 2 in \mathbb{P}^3 might contain 0, 1 or 2 twistor fibers and if a surface of degree 2 contains more than 2 twistor fibers, then it contains infinitely many of them [13]. This is completely analogous to what we found in [4] for surfaces of bidegree $(1, 1)$ in \mathbb{F} . On the contrary, for $d = 2$ we observe a difference between the two cases. Indeed, for degree 3 surfaces in \mathbb{P}^3 , the maximum number of twistor fibers contained in a smooth surface is 5 [5], while we prove here that this number reduces to 4 for smooth surfaces of bidegree $(1, 2)$ in \mathbb{F} .

In order to present our results, we need to introduce some definitions. The flag threefold can be explicitly defined as

$$\mathbb{F} := \{(p, \ell) \in \mathbb{P}^2 \times \mathbb{P}^2 \mid p\ell = 0\},$$

where $p = [p_0 : p_1, p_2], \ell = [\ell_0 : \ell_1 : \ell_2] \in \mathbb{P}^2$ and $p\ell = p_0\ell_0 + p_1\ell_1 + p_2\ell_2$. The notation (p, ℓ) , would recall a couple *(point, line)*, and the condition $p\ell = 0$ translates as *p belongs to ℓ*. In order to simplify the notation, we identify the second factor \mathbb{P}^{2v} with \mathbb{P}^2 .

We have three projection maps: $\pi_1, \pi_2, \pi : \mathbb{F} \rightarrow \mathbb{P}^2$, defined as

$$\pi_1(p, \ell) = p, \quad \pi_2(p, \ell) = \ell,$$

$$\pi(p, \ell) = \bar{p} \times \ell = [\bar{p}_1\ell_2 - \bar{p}_2\ell_1, \bar{p}_2\ell_0 - \bar{p}_0\ell_2, \bar{p}_0\ell_1 + \bar{p}_1\ell_0],$$

where the third one is the twistor map. The fibers of such three maps are the object of our investigation. In particular, for any $q \in \mathbb{P}^2$, the three fibers $\pi_1^{-1}(q), \pi_2^{-1}(q)$ and $\pi^{-1}(q)$ are curves of bidegree $(0, 1), (1, 0)$ and $(1, 1)$, respectively. While the fibers of π_1 and π_2 exhaust the family of bidegrees $(1, 0)$ and $(0, 1)$ curves, twistor fibers are a (non-open Zariski dense) subset of those of bidegree $(1, 1)$. It was shown in [4, Section 3.1] that any bidegree $(1, 1)$ curve

has Hilbert polynomial equal to $2t + 1$ with respect to the standard Segre embedding of $\mathbb{P}^2 \times \mathbb{P}^2$; for this reason we will call them *conics*. There are only two types of bidegree $(1, 1)$ curves: the reducible ones (union of a bidegree $(1, 0)$ and of a bidegree $(0, 1)$ curves intersecting at a point), and the smooth ones. All of them can be described as

$$L_{q,m} := \{(p, \ell) \in \mathbb{F} \mid pm = 0, q\ell = 0\},$$

where $q, m \in \mathbb{P}^2$ and, the reducible and smooth cases are obtained for $qm = 0$ or $qm \neq 0$, respectively.

In twistor theory, an important role is played by an antiholomorphic involution without fixed points, which identifies twistor fibers [6, 11]. In our case, this map can be defined as $j : \mathbb{F}^2 \rightarrow \mathbb{F}^2$, where

$$j(p, \ell) = (\bar{\ell}, \bar{p}).$$

A smooth conic $L_{q,m}$ is a twistor fiber if and only if $j(L_{q,m}) = L_{q,m}$ if and only if $m = \bar{q}$. Moreover, it is natural to classify objects in \mathbb{F} up to projective automorphisms coming from the lift via π of a unitary automorphism of \mathbb{P}^2 , i.e. a holomorphic isometry with respect to the Fubini-Study metric. Such transformations of \mathbb{F} are exactly those which commute with j (see [4, Lemma 5.4] for the flag manifold case). Thus, in particular, the number of twistor fibers contained in a given surface and their arrangement are unitary invariants.

In order to state our main results we need some more notation. We denote by $\mathcal{C}(1) = \mathcal{C}$ the set of smooth conics in \mathbb{F} and by $\mathcal{C}(n)$, $n \geq 2$, the set of n pairwise disjoint smooth conics. In an analogous way we define $\mathcal{T}(1) = \mathcal{T} \subset \mathcal{C}$ as the set of twistor fibers and $\mathcal{T}(n) \subset \mathcal{C}(n)$, $n \geq 2$, as the set of n pairwise disjoint twistor fibers.

We will see in Remark 2.5 that for any couple of different smooth conics, there is a unique bidegree $(1, 0)$ curve $L = \pi_2^{-1}(q_2)$ and a unique bidegree $(0, 1)$ curve $R = \pi_1^{-1}(q_1)$ such that L and R intersect both smooth conics. In the case of a couple of twistor fibers we also have $R = j(L)$. We say that three or more smooth conics are *collinear* if there is a $(1, 0)$ curve L which intersects all of them. To be collinear, for three or more smooth conics, is a Zariski closed condition.

To be more precise, in Definition 2.7, we define the set $\mathcal{C}^*(n)$ which parametrizes all $A \in \mathcal{C}(n)$ such that $\#(L \cap A) \leq 2$ for all curves L of bidegree $(1, 0)$. Clearly $\mathcal{C}^*(1) = \mathcal{C}(1)$, and $\mathcal{C}^*(2) = \mathcal{C}(2)$, while for $n \geq 3$ the open set $\mathcal{C}^*(n)$ is given by the set of disjoint smooth conics such that no three of them are collinear. Moreover, we set $\mathcal{T}^*(n) := \mathcal{T}(n) \cap \mathcal{C}^*(n)$. In Theorem 2.19 we characterize the elements $A \in \mathcal{C}^*(d + 1)$ to be those which do not obstruct the linear system $|\mathcal{J}_A(1, d)|$.

We now summarize the main results of the paper. In Section 3, we study surfaces of bidegree $(1, d)$ containing a certain number of smooth conics or twistor fibers, and we prove the following two theorems.

Theorem 1.1. *For any $d \in \mathbb{N}$ and $A \in \mathcal{T}^*(d+2)$, there is no irreducible surface of bidegree $(1, d)$ containing A .*

Theorem 1.2. *Fix integer $d \geq 1$ and $0 \leq n \leq d+2$. There is an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, d)|$ containing exactly n twistor fibers.*

We also show, in Theorem 3.4, that the first result is sharp. Indeed, for any $A \in \mathcal{T}^*(n)$, with $0 \leq n \leq d+1$, we are able to prove the existence of an irreducible surface of bidegree $(1, d)$ containing A and no other twistor fibers. This last issue requires some effort and the proof is divided in a detailed analysis of several cases.

Theorem 1.2 is a consequence of Theorem 3.4 and Theorem 3.8. More precisely, in Theorem 3.4, for $0 \leq n \leq d+1$, we prove that, fixed any union A of $0 \leq n \leq d+1$ non-three-by-three collinear twistor fibers, there is an irreducible $(1, d)$ -surface containing A and no other twistor fibers. The extremal case $n = d+2$ is considered in Theorem 3.8, where we prove that, given $d+2$ general collinear twistor fibers, there is an irreducible surface of bidegree $(1, d)$ containing them.

In Section 4, we focus on surfaces of bidegree $(1, 2)$ and $(1, 3)$. The main results are summarized by the following statements:

Theorem 1.3. *Fix $0 \leq n \leq 3$. There is a smooth $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ containing exactly n twistor fibers. Moreover, there exists a bidegree $(1, 2)$ irreducible surface containing exactly 4 twistor fibers.*

Theorem 1.4. *There is no irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ containing at least 5 twistor fibers.*

Theorem 1.5. *There is no irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, 3)|$ containing at least 6 twistor fibers.*

The first existence result (Theorem 1.3) follows from Theorem 4.6, for $0 \leq n \leq 3$, and Theorem 4.10, for the case $n = 4$. In the extremal case $n = 4$, we will also show that the surfaces are singular along a line.

The two non-existence results (Theorems 1.4 and 1.5) are proved in the last Section 4.2. An essential tool is Lemma 4.11 which states that if a surface of bidegree $(1, d)$ contains $d+3$ or more collinear twistor fibers, then this surface is reducible and one of its components is a surface of bidegree $(1, 1)$ containing 4 of the prescribed twistor fibers.

The proofs of our results are based on cohomological methods and on a careful description of the linear systems of surfaces in the flag threefold.

2. Preliminaries and first results

In this section we collect some known results about algebraic curves and surfaces in the flag threefold. Then we give first results on the space of bidegree $(0, d)$ and $(1, d)$ -surfaces containing a certain number of twisted fibers. In particular, we introduce the concept of *collinear* smooth conics and give a topological characterization in terms of the cohomology of certain ideal sheaves.

For most of the known material on \mathbb{F} we refer to [4] and to [3, Section 2]. However, in order to be as self-contained as possible, we will recall some basic ideas and results here.

Consider the multi projective space $\mathbb{P}^2 \times \mathbb{P}^2$; an element $(p, \ell) \in \mathbb{P}^2 \times \mathbb{P}^2$ will be a pair written in the following form $p = [p_0 : p_1 : p_2], \ell = [\ell_0 : \ell_1 : \ell_2]^T$, so that $p\ell = p_0\ell_0 + p_1\ell_1 + p_2\ell_2$. Even though it is classically embedded in $\mathbb{P}^2 \times \mathbb{P}^{2\vee}$, we can see $\mathbb{F} := \{(p, \ell) \in \mathbb{P}^2 \times \mathbb{P}^2 \mid p\ell = 0\}$ as a hypersurface of bidegree $(1, 1)$ of $\mathbb{P}^2 \times \mathbb{P}^2$. We denote by Π_1 and Π_2 the two standard projections of $\mathbb{P}^2 \times \mathbb{P}^2$ and use lower case for their restrictions, i.e. $\pi_i = \Pi_i|_{\mathbb{F}}, i = 1, 2$. Thus, the two natural projections define a natural notion of bidegree for algebraic surfaces in \mathbb{F} . Furthermore, for all $(a, b) \in \mathbb{Z}^2$ we have the following natural exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^2}(a - 1, b - 1) \rightarrow \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^2}(a, b) \rightarrow \mathcal{O}_{\mathbb{F}}(a, b) \rightarrow 0, \tag{1}$$

and, for any $(a, b) \in \mathbb{N}^2$, we get (see e.g. [4, Lemma 2.3])

$$h^0(\mathcal{O}_{\mathbb{F}}(a, b)) = \frac{(a + 1)(b + 1)(a + b + 2)}{2} \quad \text{and} \quad h^1(\mathcal{O}_{\mathbb{F}}(a, b)) = 0. \tag{2}$$

It will be useful to recall from [4, Proposition 3.11] the multiplication rules in the Chow ring:

$$\begin{aligned} \mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(1, 0) &= 0, & \mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(0, 1) \cdot \mathcal{O}_{\mathbb{F}}(1, 0) &= 1, \\ \mathcal{O}_{\mathbb{F}}(0, 1) \cdot \mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(0, 1) &= 1, & \mathcal{O}_{\mathbb{F}}(0, 1) \cdot \mathcal{O}_{\mathbb{F}}(0, 1) \cdot \mathcal{O}_{\mathbb{F}}(0, 1) &= 0. \end{aligned} \tag{3}$$

2.1. Curves in \mathbb{F} and smooth conics. Let us recall the notion of bidegree for the family of algebraic curves in \mathbb{F} already given in [4, 3].

Definition 2.1. Let $C \subset \mathbb{F}$ be an irreducible and reduced algebraic curve. We define the bidegree of C as the couple of positive integers (d_1, d_2) , where $d_i = 0$ if $\pi_i(C) = \{x\}$, otherwise $d_i = \deg(\pi_i(C)) \deg(\pi_i|_C)$.

If a curve D has irreducible components C_1, \dots, C_s then the bidegree of D is the sum of the bidegrees of C_1, \dots, C_s .

Recall from [3, Remark 2.4] that if a curve C is such that $C \cdot \mathcal{O}_{\mathbb{F}}(1, 0) = d_1$ and $C \cdot \mathcal{O}_{\mathbb{F}}(0, 1) = d_2$, then it has bidegree (d_1, d_2) .

From the multiplication table (3) we can easily derive the following formula.

Lemma 2.2. For any choice of non-negative integers a, b, c, d , the one-dimensional cycle $\mathcal{O}_{\mathbb{F}}(a, b) \cdot \mathcal{O}_{\mathbb{F}}(c, d)$ has bidegree

$$(ad + b(c + d), a(c + d) + bc).$$

Proof. We have $\mathcal{O}_{\mathbb{F}}(a, b) \cdot \mathcal{O}_{\mathbb{F}}(c, d) = ac\mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(1, 0) + (ad + bc)\mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(0, 1) + bd\mathcal{O}_{\mathbb{F}}(0, 1) \cdot \mathcal{O}_{\mathbb{F}}(0, 1)$. So the thesis is easily obtained by recalling that $\mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(1, 0)$ (resp. $\mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(0, 1)$, resp. $\mathcal{O}_{\mathbb{F}}(0, 1) \cdot \mathcal{O}_{\mathbb{F}}(0, 1)$) is a one-dimensional cycle of bidegree $(0, 1)$ (resp. bidegree $(1, 1)$, resp. bidegree $(1, 0)$). \square




FIGURE 1. Any two smooth conics are connected by a curve of bidegree (1, 0) and by a curve of (0, 1).

Remark 2.3. Note that the fibers of π_1 are algebraic curves of bidegree (0, 1), while those of π_2 have bidegree (1, 0) (see, e.g. [4, Section 3]). Moreover, all bidegree (0, 1) curves can be seen as complete intersections of two different (1, 0)-surfaces (and analogously for bidegree (1, 0) curves).

Among all the algebraic curves in \mathbb{F} , we focus our attention on the family of bidegree (1, 1) curves. These are described geometrically in [4, Section 3.1] and are parameterized by $(q, m) \in \mathbb{P}^2 \times \mathbb{P}^2$. In fact, as anticipated in the introduction, each of these curves can be written as

$$L_{q,m} := \{(p, \ell) \in \mathbb{F} \mid p \in m, \ell \ni q\} = \{(p, \ell) \in \mathbb{F} \mid q\ell = 0, pm = 0\}.$$

There are two types of these curves: the smooth and irreducible ones (when $qm \neq 0$) and the union of a (1, 0) and of a (0, 1) intersecting at one point (when $qm = 0$, i.e. $(q, m) \in \mathbb{F}$). In any case, any curve of bidegree (1, 1) can be seen as the complete intersection of a surface of bidegree (1, 0) with one of bidegree (0, 1). As already mentioned in the introduction, the 4-dimensional family of smooth irreducible (1, 1) curves is denoted by \mathcal{C} . The elements of \mathcal{C} are called *smooth conics*.

Remark 2.4. From the definition of smooth conics, it is clear that, for any $C \in \mathcal{C}$ we have that $\pi_i(C)$ is a line in \mathbb{P}^2 .

Remark 2.5. Note that for any two different elements $L_{q,m}, L_{q',m'} \in \mathcal{C}$ there exists a unique curve L of bidegree (1, 0) and a unique curve R of bidegree (0, 1) such that L and R meet both $L_{q,m}$ and $L_{q',m'}$ at one point (see Figure 1). From the analysis made in [4, Section 3.1] it is easy to see that $L = \pi_2^{-1}(q \times q')$ and $R = \pi_1^{-1}(m \times m')$, where \times stands for the standard (formal) cross product. Equivalently, $L = \pi_2^{-1}(\text{Sing}(\pi_2(A)))$ and $R = \pi_1^{-1}(\text{Sing}(\pi_1(A)))$. We say that three or more disjoint smooth conics are *collinear* if they intersect the same (1, 0) curve L .

The fibers of the twistor projection $\pi : \mathbb{F} \rightarrow \mathbb{P}^2$ (see [4, Section 5]) form a subset \mathcal{T} of the family of conics \mathcal{C} . The twistor fibers are also characterized to be the irreducible elements in \mathcal{C} that are fixed by the anti-holomorphic involution $j : \mathbb{F} \rightarrow \mathbb{F}$ defined as

$$j(p, \ell) = (\bar{\ell}, \bar{p}).$$

Being the set of fixed points of j , a curve $L_{q,m}$ belongs to \mathcal{T} if and only if $m = \bar{q}$. Furthermore, the set \mathcal{T} is Zariski dense in \mathcal{C} (see, e.g. [3, Section 4]).

Remark 2.6. If L is the bidegree curve $(1, 0)$ connecting two different twistor fibers (see Remark 2.5), then the curve of bidegree $(0, 1)$ connecting them is exactly $R = j(L)$. Thus, if three twistor fibers are collinear, then they intersect the same $(1, 0)$ curve and the same $(0, 1)$ curve.

Recall from the introduction that, for any positive integer n , $\mathcal{C}(n)$ denotes the $4n$ -dimensional set of n pairwise disjoint elements of \mathcal{C} and $\mathcal{T}(n)$ denotes the set of n pairwise disjoint elements of \mathcal{T} . As before, $\mathcal{T}(n)$ is Zariski dense in $\mathcal{C}(n)$ (see again [3, Section 4]).

We now introduce the following crucial definition.

Definition 2.7. For any $n \geq 1$ let $\mathcal{C}^*(n)$ be the set of all $A \in \mathcal{C}(n)$ such that for any curve L of bidegree $(1, 0)$, it holds $\#(L \cap A) \leq 2$.

Set $\mathcal{T}^*(n) := \mathcal{T}(n) \cap \mathcal{C}^*(n)$.

Obviously we have $\mathcal{C}^*(n) = \mathcal{C}(n)$ and $\mathcal{T}^*(n) = \mathcal{T}(n)$, for $n = 1, 2$. For $n \geq 3$ the set $\mathcal{C}(n) \setminus \mathcal{C}^*(n)$ parameterizes unions of n disjoint smooth conics such that at least three of them are collinear, hence $\mathcal{C}^*(n)$ is open and Zariski dense in $\mathcal{C}(n)$, as well as $\mathcal{T}^*(n)$ in $\mathcal{T}(n)$. Therefore for any $n \geq 1$, all of the following inclusions are Zariski dense: $\mathcal{T}^*(n) \subset \mathcal{T}(n) \subset \mathcal{C}(n)$, $\mathcal{T}^*(n) \subset \mathcal{C}^*(n) \subset \mathcal{C}(n)$.

2.2. Surfaces of bidegree $(1, 0)$ and $(0, 1)$. Now we turn our attention back to surfaces. We recall from [4, Section 3.2] and [3, Section 2] that $(1, 0)$ and $(0, 1)$ -surfaces are Hirzebruch surfaces of the first type. In particular, a surface X of bidegree $(1, 0)$ can be seen as the lift, via π_1 , of a line (and analogously for a surface Y of bidegree $(0, 1)$). Using this description, it is easy to see that each of these surfaces represents the blow up of \mathbb{P}^2 at one point. Let F_1 be a Hirzebruch surface of type 1; we now describe the relation between the generators of the Picard group of F_1 and the family of curves in \mathbb{F} described earlier. We recall that $\text{Pic}(F_1) = \mathbb{Z}h \oplus \mathbb{Z}f$, where

$$h^2 = -1, \quad f^2 = 0, \quad hf = 1.$$

Notation 2.8. For the following analysis and the rest of the paper, X will denote a surface of bidegree $(1, 0)$, while Y a surface of bidegree $(0, 1)$.

If we identify a surface X with F_1 , we get that $\mathcal{O}_X(1, 0) \simeq \mathcal{O}_{F_1}(f)$, which again corresponds to the set of curves in \mathbb{F} of bidegree $(0, 1)$ contained in X . On the other hand, we have that $\mathcal{O}_X(0, 1) \simeq \mathcal{O}_{F_1}(h + f)$, which corresponds to the elements of \mathcal{C} .

Thus, for any $a, b \in \mathbb{Z}$ and for any $\alpha, \beta \in \mathbb{Z}$, we obtain the following two relations

$$\mathcal{O}_X(a, b) \cong \mathcal{O}_{F_1}(bh + (a + b)f), \quad \text{and} \quad \mathcal{O}_{F_1}(\alpha h + \beta f) \cong \mathcal{O}_X(\beta - \alpha, \alpha). \quad (4)$$

For a surface Y of bidegree $(0, 1)$ we can derive similar formulæ:

$$\mathcal{O}_Y(a, b) \cong \mathcal{O}_{F_1}(ah + (a + b)f), \quad \text{and} \quad \mathcal{O}_{F_1}(\alpha h + \beta f) \cong \mathcal{O}_Y(\alpha, \beta - \alpha), \quad (5)$$

for any $a, b \in \mathbb{Z}$ and for any $\alpha, \beta \in \mathbb{Z}$.

Remark 2.9. Let X be a surface of bidegree $(1, 0)$. Then X contains no element of $\mathcal{C}(2)$. In fact, every element of \mathcal{C} in X corresponds to an element of $\mathcal{O}_{F_1}(h + f)$ and any two elements of type $h + f$ meet. The same holds for surfaces Y of bidegree $(0, 1)$. In particular, for every surface of bidegree $(1, 0)$ or $(0, 1)$ there is exactly one twistor fiber contained in it.

We now recall from [3, Lemma 2.5] that, for any $a, b \geq 0$, by using the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{F}}(a - 1, b) \rightarrow \mathcal{O}_{\mathbb{F}}(a, b) \rightarrow \mathcal{O}_X(a, b) \rightarrow 0,$$

and its analog for Y , we have that

$$h^0(\mathcal{O}_X(a, b)) = a(b + 1) + \binom{b + 2}{2}, \quad h^0(\mathcal{O}_Y(a, b)) = b(a + 1) + \binom{a + 2}{2},$$

while

$$h^1(\mathcal{O}_X(a, b)) = h^1(\mathcal{O}_Y(a, b)) = 0.$$

Furthermore, if $a > 0, b > 0$, the line bundles $\mathcal{O}_X(a, b)$ and $\mathcal{O}_Y(a, b)$ are very ample.

2.3. Surfaces of bidegree $(0, d)$ and $(1, d)$. We now move on to studying higher bidegree surfaces. We start with some considerations about bidegree $(0, d)$ surfaces.

Remark 2.10. As described in [4, Section 3.3], every irreducible and reduced surface S of bidegree $(0, d)$ is equal to $\pi_1^{-1}(C)$ for some irreducible and reduced curve C of degree d . Therefore, for $d \geq 2$, no irreducible and reduced $S \in |\mathcal{O}_{\mathbb{F}}(0, d)|$ contains a smooth conic. Otherwise, thanks to Remark 2.4, $\pi_i(S)$ would contain a line, but $\pi_1(S)$ is an irreducible and reduced curve of degree d .

For $n \geq 2$ we now compute how many bidegree $(0, d)$ -surfaces contain a fixed element of $\mathcal{C}(n)$. First we set up the following notation.

Notation 2.11. We will denote by $\mathcal{J}_{U, V}$ the ideal sheaf of a scheme U contained in a projective variety V ; whenever $V = \mathbb{F}$ we will omit it. In particular, if $A \in \mathcal{C}(n)$ we will write $\mathcal{J}_A := \mathcal{J}_{A, \mathbb{F}}$.

Lemma 2.12. Fix $d \geq 0, n \geq 1$, and $A \in \mathcal{C}(n)$. We have

$$h^0(\mathcal{J}_A(0, d)) = \frac{(d - n + 2)(d - n + 1)}{2}$$

and

$$h^1(\mathcal{J}_A(0, d)) = \begin{cases} \frac{n(n-1)}{2} & \text{if } n \leq d + 1 \\ n(d + 1) - \frac{(d+2)(d+1)}{2} & \text{if } n \geq d + 1. \end{cases}$$

Proof. Recall that $\mathcal{O}_{\mathbb{F}}(0, d) = \pi_1^*(\mathcal{O}_{\mathbb{P}^2}(d))$, and $\mathcal{J}_A(0, d) = \pi_1^*(\mathcal{J}_{T, \mathbb{P}^2}(d))$ where $T = \pi_1(A)$ is a union of n distinct lines in \mathbb{P}^2 . In general, we have that:

$$h^0(\mathcal{O}_{\mathbb{F}}(0, d)) = \binom{d+2}{2}, \quad h^0(\mathcal{J}_A(0, d)) = \binom{d+2-n}{2},$$

$$h^0(\mathcal{O}_A(0, d)) = n(d+1),$$

hence, using the exact sequence

$$0 \rightarrow \mathcal{J}_A(0, d) \rightarrow \mathcal{O}_{\mathbb{F}}(0, d) \rightarrow \mathcal{O}_A(0, d) \rightarrow 0,$$

since $h^1(\mathcal{O}_{\mathbb{F}}(0, d)) = 0$ and $\binom{d+2-n}{2} = 0$ if $n \geq d + 1$, we get the result. \square

Remark 2.13. As a direct consequence of the previous lemma, we can state that, for any $C \in \mathcal{C}$, there is only one surface Y in $|\mathcal{J}_C(0, 1)|$ and, analogously, only one surface X in $|\mathcal{J}_C(1, 0)|$.

Remark 2.14. If $A \in \mathcal{C}(n)$, we consider the following exact sequence:

$$0 \rightarrow \mathcal{J}_A(a, b) \rightarrow \mathcal{O}_{\mathbb{F}}(a, b) \rightarrow \mathcal{O}_A(a, b) \rightarrow 0.$$

Since the conics in A are all disjoint, for each $(a, b) \in \mathbb{N}^2$ we have that

$$h^0(\mathcal{O}_A(a, b)) = n(a + b + 1), \tag{6}$$

(see, e.g. [3, Sequence (7) and proof of Theorem 4.4]). Recall from Formula (2) that

$$h^0(\mathcal{O}_{\mathbb{F}}(a, b)) = \frac{(a+1)(b+1)(a+b+2)}{2}.$$

Therefore, as $h^1(\mathcal{O}_A(a, b)) = h^1(\mathcal{O}_{\mathbb{F}}(a, b)) = 0$, for any $A \in \mathcal{C}(n)$ and $d \geq 0$, we have

$$\chi(\mathcal{J}_A(1, d)) = h^0(\mathcal{J}_A(1, d)) - h^1(\mathcal{J}_A(1, d)) = (d+1)(d+3) - n(d+2). \tag{7}$$

Hence, if $n = d + 2$, we have $h^0(\mathcal{O}_A(1, d)) = (d + 2)^2$ and so

$$h^0(\mathcal{O}_{\mathbb{F}}(1, d)) = (d+1)(d+3) = (d+2)^2 - 1 = h^0(\mathcal{O}_A(1, d)) - 1 \tag{8}$$

and, if $n = d + 1$:

$$h^0(\mathcal{O}_{\mathbb{F}}(1, d)) = h^0(\mathcal{O}_A(1, d)) + d + 1.$$

In what follows, we are going to implicitly use the following simple observation.

Remark 2.15. Since the elements of $\mathcal{J}(1)$ are the fibers of the twistor map, each $p \in \mathbb{F}$ is contained in a unique element, C , of \mathcal{J} . Since $j(C) = C$, $j(p) \in C$. Analogously, note that p is contained in a unique curve of bidegree $(1, 0)$ and a unique curve of bidegree $(0, 1)$, $\pi_2^{-1}(\pi_2(p))$ and $\pi_1^{-1}(\pi_1(p))$.

The following remark will be used several times in the next pages.

Remark 2.16. Fix a positive integer d and an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, d)|$. Since $\pi_{1|S}$ is birational onto its image, S is rational. By the Bézout Theorem, for any $p \in \mathbb{P}^2$, the bidegree $(1, 0)$ curve $\pi_2^{-1}(p)$ is either contained in S or intersects S in a single point (scheme-theoretically).

We conclude this subsection with a technical result that will be used in the next pages. In [3, Proposition 4.1] we proved that there exists at most one surface of bidegree (a, b) containing a number equal to or greater than $a^2 + ab + b^2$ of smooth conics. We will now generalize this result to a more general context.

Proposition 2.17. Fix $(a, b, c, d) \in \mathbb{N}^4$ so that $(a, b) \neq (0, 0)$, $c > 0$, $d > 0$. Take $A \in \mathcal{C}(n)$. Assume the existence of an irreducible and reduced $S \in |\mathcal{O}_{\mathbb{F}}(a, b)|$ containing A and assume one of the following conditions:

- (1) $ad + b(c + d) < n$;
- (2) $a(c + d) + bc < n$;
- (3) $ad + b(c + d) = a(c + d) + bc = n$.

Then every element of $|\mathcal{J}_A(c, d)|$ contains S and in particular $c \geq a$ and $d \geq b$.

Proof. Assume by contradiction the existence of $S' \in |\mathcal{J}_A(c, d)|$ such that $S' \not\supseteq S$. We have

$$\begin{aligned} \mathcal{O}_{\mathbb{F}}(a, b) \cdot \mathcal{O}_{\mathbb{F}}(c, d) &= ac\mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(1, 0) + (ad + bc)\mathcal{O}_{\mathbb{F}}(1, 0) \cdot \mathcal{O}_{\mathbb{F}}(0, 1) \\ &\quad + bd\mathcal{O}_{\mathbb{F}}(0, 1) \cdot \mathcal{O}_{\mathbb{F}}(0, 1). \end{aligned}$$

Since $S' \not\supseteq S$, the intersection $S \cap S'$ is a curve of bidegree $(ad + b(c + d), a(c + d) + bc)$ (see Lemma 2.2). Since $c > 0$ and $d > 0$, then $\mathcal{O}_{\mathbb{F}}(c, d)$ is ample. Moreover, since S is irreducible, A has bidegree (n, n) , and A is not connected, then the intersection contains some more components in addition to A . So either $ad + bc + bd > n$ and $ac + ad + bc \geq n$ or $ad + bc + bd \geq n$ and $ac + ad + bc > n$. \square

2.4. Non-collinear smooth conics. We now want to characterize the conics in $\mathcal{C}^*(n)$ in terms of cohomology. We start by showing that the vanishing of certain cohomology groups implies that an element $A \in \mathcal{C}(n)$ lies in $\mathcal{C}^*(n)$.

Lemma 2.18. Fix $d \geq 0$, $3 \leq n \leq d + 1$ and $A \in \mathcal{C}(n)$. If $h^1(\mathcal{J}_A(1, d)) = 0$ then $A \in \mathcal{C}^*(n)$.

Proof. Suppose, by contradiction, that there exists a curve L of bidegree $(1, 0)$ such that $\#(L \cap A) \geq 3$ and consider the exact sequence that defines \mathcal{J}_A :

$$0 \rightarrow \mathcal{J}_A(1, d) \rightarrow \mathcal{O}_{\mathbb{F}}(1, d) \rightarrow \mathcal{O}_A(1, d) \rightarrow 0. \tag{9}$$

We will prove that the restriction map $H^0(\mathcal{O}_{\mathbb{F}}(1, d)) \rightarrow H^0(\mathcal{O}_A(1, d))$ is not surjective, which implies that $h^1(\mathcal{J}_A(1, d)) > 0$. Let us assume that it is surjective.

Then consider the following diagram

$$\begin{array}{ccc} \mathcal{O}_{\mathbb{F}}(1, d) & \longrightarrow & \mathcal{O}_A(1, d) \\ \downarrow & & \downarrow \\ \mathcal{O}_L(1, d) & \longrightarrow & \mathcal{O}_{L \cap A}(1, d). \end{array}$$

Since the vertical maps are surjective, the induced map

$$H^0(\mathcal{O}_{\mathbb{F}}(1, d)) \rightarrow H^0(\mathcal{O}_{A \cap L}(1, d))$$

is also surjective and thus of rank at least 3 (since $L \cap A$ has cardinality at least 3 and the irreducible components of A are pairwise disjoint). On the other hand, the restriction map $H^0(\mathcal{O}_{\mathbb{F}}(1, d)) \rightarrow H^0(\mathcal{O}_L(1, d))$ has rank 2, and this leads to a contradiction. \square

Before completing the characterization of conics in $\mathcal{C}^*(d + 1)$, we will introduce a general construction that will be used in several subsequent discussions. Let $A \in \mathcal{C}(n)$ and let C be any connected component of A . Set $B := A \setminus C$. Then, for any $a, b \geq 0$, if $Y \in |\mathcal{O}_C(0, 1)|$, we have the following residual exact sequence

$$0 \rightarrow \mathcal{J}_{Res_Y(A)}(a, b - 1) \rightarrow \mathcal{J}_A(a, b) \rightarrow \mathcal{J}_{A \cap Y}(a, b) \rightarrow 0,$$

but since $Res_Y(A) = B$ and $A \cap Y = (B \cap Y) \cup C$, we have

$$0 \rightarrow \mathcal{J}_B(a, b - 1) \rightarrow \mathcal{J}_A(a, b) \rightarrow \mathcal{J}_{(B \cap Y) \cup C}(a, b) \rightarrow 0. \tag{10}$$

Obviously, an analogous sequence can be written for $X \in |\mathcal{O}_C(1, 0)|$.

Theorem 2.19. *Fix $d \geq 0$ and $A \in \mathcal{C}(d + 1)$. We have $A \in \mathcal{C}^*(d + 1)$ if and only if $h^1(\mathcal{J}_A(1, d)) = 0$.*

Proof. Thanks to the previous lemma we only need to prove that $A \in \mathcal{C}^*(d + 1)$ satisfies $h^1(\mathcal{J}_A(1, d)) = 0$. We use induction on $d \geq 0$. The case $d = 0$ is true by Lemma 2.12. So we can assume $d > 0$ and use induction on d . Let C be a connected component of A , set $B := A \setminus C$ and call Y the unique element of $|\mathcal{J}_C(0, 1)|$. Consider the residual exact sequence (10), with $a = 1$ and $b = d$. Since $A \in \mathcal{C}(d + 1)$, $C \cap B = \emptyset$ and $B \cap Y$ is a set of d different points, up to the identification of D with F_1 we have

$$\mathcal{J}_{(B \cap Y) \cup C}(1, d) \cong \mathcal{J}_{(B \cap Y) \cup C, F_1}(1, d)(h + (d + 1)f) \cong \mathcal{J}_{B \cap Y, F_1}(df).$$

Using (10) and induction, we are left to prove that $h^1(\mathcal{J}_{B \cap Y, F_1}(df)) = 0$ if and only if $A \in \mathcal{C}^*(d + 1)$.

Consider now the following exact sequence

$$0 \rightarrow \mathcal{J}_{B \cap Y, F_1}(df) \rightarrow \mathcal{O}_{F_1}(df) \rightarrow \mathcal{O}_{B \cap Y}(df) \rightarrow 0. \tag{11}$$

Since $h^0(\mathcal{O}_{F_1}(df)) = d + 1$ and $h^0(\mathcal{O}_{B \cap Y}(df)) = d$, we have the following inequality $h^1(\mathcal{J}_{B \cap Y, F_1}(df)) > 0$ if and only if $h^0(\mathcal{J}_{B \cap Y, F_1}(df)) \geq 2$. The last inequality means that there are at least two different sets of d fibers containing the set of d points $B \cap Y$. This is equivalent to the fact that there exists a fiber

$L \in |f|$ such that $\#(B \cap L) \geq 2$. Since L is a curve of bidegree $(1, 0)$ in \mathbb{F} , then $L \cdot Y = 0$ in the intersection ring of \mathbb{F} , so $L \subset Y$, and we get $L \cap C \neq \emptyset$. Thus $\#(L \cap A) \geq 3$, which means that $A \notin \mathcal{C}^*(d + 1)$. \square

Corollary 2.20. *Fix $d \geq 0, 0 \leq n \leq d + 1$ and $A \in \mathcal{C}^*(n)$. Then $h^1(\mathcal{J}_A(1, d)) = 0$. In particular, for $n = 0, 1, 2$ and for any $A \in \mathcal{C}(n)$, we have*

$$h^1(\mathcal{J}_A(1, 1)) = 0 \quad \text{and} \quad h^0(\mathcal{J}_A(1, 1)) = 8 - 3n.$$

Proof. By using [3, Remark 4.3], we easily get the first part of the statement. The second part follows from Formula (7). \square

We note that since $\mathcal{T}(n)$ is Zariski dense in $\mathcal{C}(n)$, then the characterization given by Theorem 2.19 also holds for the set $\mathcal{T}^*(n)$.

Lemma 2.21. *Fix an integer $d \geq 0$ and $A \in \mathcal{C}^*(d + 2)$. Then $h^1(\mathcal{J}_A(1, d)) \leq 1$.*

Proof. The lemma is true for $d = 0$, because $h^0(\mathcal{J}_A(1, 0)) = 0$ (see Remark 2.9). We assume $d > 0$ and use induction on d . Let C be a connected component of A , set $B := A \setminus C$ and call Y the unique element of $|\mathcal{J}_C(0, 1)|$. Consider the residual exact sequence (10), with $a = 1$ and $b = d$. Since $A \in \mathcal{C}^*(d + 2)$, $C \cap B = \emptyset$ and $B \cap Y$ is formed by $d + 1$ different points, up to the identification of D with F_1 we have

$$\mathcal{J}_{(B \cap Y) \cup C, Y}(1, d) \cong \mathcal{J}_{(B \cap Y) \cup C, F_1}(1, d)(h + (d + 1)f) \cong \mathcal{J}_{B \cap Y, F_1}(df).$$

Using (10) and induction, we are left to prove that $h^1(\mathcal{J}_{B \cap Y, F_1}(df)) = 0$ if $A \in \mathcal{C}^*(d + 2)$.

Consider now the following exact sequence

$$0 \rightarrow \mathcal{J}_{B \cap Y, F_1}(df) \rightarrow \mathcal{O}_{F_1}(df) \rightarrow \mathcal{O}_{B \cap Y}(df) \rightarrow 0. \tag{12}$$

Since

$$h^0(\mathcal{O}_{F_1}(df)) = d + 1, \quad h^0(\mathcal{O}_{B \cap Y}(df)) = d + 1,$$

we have that $h^1(\mathcal{J}_{B \cap Y, F_1}(df)) > 0$ if and only if $h^0(\mathcal{J}_{B \cap Y, F_1}(df)) > 0$. This is equivalent to the fact that there exists a fiber $L \in |f|$ such that $\#(B \cap L) \geq 2$. Since L is a curve of bidegree $(1, 0)$ in \mathbb{F} , then $L \cdot Y = 0$ in the intersection ring of \mathbb{F} , therefore $L \subset Y$, and so we get $L \cap C \neq \emptyset$. Thus $\#(L \cap A) \geq 3$, which means that $A \notin \mathcal{C}^*(d + 2)$. \square

As said in Remark 2.5, for any element $A \in \mathcal{C}(2)$ there exists a unique curve L of bidegree $(1, 0)$ and a unique curve R of bidegree $(0, 1)$ such that both intersect the elements of A at a point. As described in the following result, it turns out that $A \cup L \cup R$ is the base locus of $|\mathcal{J}_A(1, 1)|$.

Proposition 2.22. *For any $A \in \mathcal{C}(2)$, we have that*

- (1) *the general element in $|\mathcal{J}_A(1, 1)|$ is irreducible;*
- (2) *the base locus \mathcal{B} of $|\mathcal{J}_A(1, 1)|$ is $A \cup L \cup R$, where L and R are the curves described in Remark 2.5.*

Proof. Since $A \in \mathcal{C}(2)$, we have $h^1(\mathcal{J}_A(1, 1)) = 0$ and $h^0(\mathcal{J}_A(1, 1)) = 2$ by Corollary 2.20. Let C_1 and C_2 be the connected components of A . Denote by X_i the only element of $|\mathcal{O}_{\mathbb{F}}(1, 0)|$ containing C_i and by Y_i the only element of $|\mathcal{O}_{\mathbb{F}}(0, 1)|$ containing C_i . The surfaces $X_1 \cup Y_2$ and $X_2 \cup Y_1$ are the only reducible elements of $|\mathcal{J}_A(1, 1)|$ and hence, the general element in $|\mathcal{J}_A(1, 1)|$ is irreducible and (1) is proved.

To prove (2) we analyze the base locus \mathcal{B} of $|\mathcal{J}_A(1, 1)|$. If $S, S' \in |\mathcal{J}_A(1, 1)|$ are irreducible and $S \neq S'$, then the one-dimensional cycle $S \cap S'$ has bidegree $(3, 3)$ and it contains A , which has bidegree $(2, 2)$. Let $R := X_1 \cap X_2$ and $L := Y_1 \cap Y_2$, where X_i and Y_i are the surfaces defined in the first part of this proof. The curves L and R are exactly those given in Remark 2.5. In particular, $\#(L \cap A) = \#(R \cap A) = 2$ and hence, by Bézout and Remark 2.16, $L \cup R \subset \mathcal{B}$. Furthermore, recall that the reducible surfaces, $X_1 \cup Y_2$ and $X_2 \cup Y_1$ belong to $|\mathcal{J}_A(1, 1)|$ and their intersection $(X_1 \cup Y_2) \cap (X_2 \cup Y_1)$ is $A \cup L \cup R$. So the base locus of $|\mathcal{J}_A(1, 1)|$ is exactly $\mathcal{B} = A \cup L \cup R$. \square

Remark 2.23. By generalizing the proof of Proposition 2.22, we can say something about the base locus of $\mathcal{J}_A(1, d)$, for $A \in \mathcal{C}(n)$. Fix the integers $d > 0$ and $n \geq 2$ and take any $A \in \mathcal{C}(n)$. Then, the base locus \mathcal{B} of $\mathcal{J}_A(1, d)$ contains all curves L of bidegree $(1, 0)$ such that $\#(L \cap A) \geq 2$. If $A \in \mathcal{C}^*(n)$, then there are exactly $\binom{n}{2}$ such curves L (the number of lines connecting two points in a set of n general points). If $A \in \mathcal{T}(n)$, then $\#(j(L) \cap A) \geq 2$ for all L such that $\#(L \cap A) \geq 2$, since $j(A) = A$.

3. Surfaces of bidegree $(1, d)$

In this section we prove Theorems 1.1 and 1.2. In particular, we give some results for the case of a surface of bidegree $(1, d)$. Later, in the following section, we will specialize to the cases $d = 2$ and $d = 3$. The case $d = 1$ has been studied in great detail in [4], and here we add a simple lemma useful for what follows.

Lemma 3.1. *For any $A \in \mathcal{T}^*(3)$, there is no irreducible surface of bidegree $(1, 1)$ containing A .*

Proof. Suppose there exists an irreducible surface M of bidegree $(1, 1)$ containing $A \in \mathcal{T}^*(3)$. First of all, thanks to [4, Corollary 8.4], we have that $M = j(M)$. Then, as explained at the beginning of Section 7.1 in [4], either M is smooth or reducible. But if M is a smooth j -invariant surface of bidegree $(1, 1)$ containing 3 twistor fibers, then it contains infinitely many of them, and these are parameterized by a circle (see [4, Theorem 7.2]). However, smooth surfaces of bidegree $(1, 1)$ can be seen as the blow up of \mathbb{P}^2 at three points both via π_1 and π_2 . In particular, up to unitary transformations, it is possible to write M as the set $\{([p_0 : p_1 : p_2], [\ell_0 : \ell_1 : \ell_2]) \in \mathbb{F} \mid p_1 \ell_1 + \lambda p_2 \ell_2\}$, with $\lambda \in \mathbb{R} \setminus \{0, 1\}$. In these coordinates, M contains $\pi_\mu^{-1}([1 : 0 : 0])$, $\pi_\mu^{-1}([0 : 1 : 0])$, $\pi_\mu^{-1}([0 : 0 : 1])$, for

$\mu = 1, 2$ and, the family of twistor fibers $\pi^{-1}([q_0 : q_1 : q_2])$ defined by:

$$\begin{cases} q_0 = 0 \text{ and } |q_1|^2\lambda + |q_2|^2 = 0 & \text{if } \lambda < 0, \\ q_1 = 0 \text{ and } |q_2|^2 - |q_0|^2(\lambda - 1) = 0 & \text{if } \lambda > 1, \\ q_2 = 0 \text{ and } |q_1|^2\lambda + |q_0|^2(\lambda - 1) = 0 & \text{if } 0 < \lambda < 1. \end{cases}$$

Take for instance $\lambda < 0$, then every twistor fiber in M intersects the line $L = \pi_2^{-1}([1 : 0 : 0])$ of bidegree $(1, 0)$. An analogous consideration holds if $0 < \lambda < 1$ or $\lambda > 1$. So we get a contradiction. \square

In the previous lemma we showed that an irreducible $(1, 1)$ -surface cannot contain three twistor fibers in general position. On the other hand if M is a $(1, 1)$ -surface containing a given $A \in \mathcal{T}(3) \setminus \mathcal{T}^*(3)$, then by [4, Corollary 8.3] we have that M is j -invariant, so it is either smooth or reducible. Moreover, if M contains infinitely many twistor fibers, then all of them intersect a bidegree $(1, 0)$ curve L and its associated $(0, 1)$ curve $R = j(L)$.

Remark 3.2. In [4, Section 8.1] we gave examples of bidegree $(1, 1)$ smooth surfaces containing exactly 0, 1 or 2 twistor fibers.

Remark 3.3. Since a smooth surface of bidegree $(1, 1)$ is a Del Pezzo surface of degree 6, it is characterized either by the three curves of bidegree $(1, 0)$ that it contains, or by the three curves of bidegree $(0, 1)$ that it contains. In fact, recall that these surfaces represent the blow up of \mathbb{P}^2 at three points with respect to either π_1 or π_2 . Note that if a smooth surface of bidegree $(1, 1)$ is j -invariant, then it is uniquely determined by three twistor fibers contained in it, and not by the curves L and $R = j(L)$ (of bidegree $(1, 0)$ and $(0, 1)$, respectively) which intersect all the twistor fibers.

Now we are ready to give the proof of our first main theorem.

Proof of Theorem 1.1: Thanks to Remark 2.9 and Lemma 3.1, the result is true for $d = 0$ and $d = 1$.

Suppose now that $d \geq 2$ and, by contradiction, that S is an irreducible $(1, d)$ -surface containing $A \in \mathcal{T}^*(d + 2)$. Call C a connected component of A and set $B := A \setminus C$. Let Y be the unique (by Lemma 3.2) element of $|\mathcal{J}_C(0, 1)|$ and consider the following exact sequence (which is a special case of the one in formula (10)):

$$0 \rightarrow \mathcal{J}_B(1, d - 1) \rightarrow \mathcal{J}_A(1, d) \rightarrow \mathcal{J}_{(B \cap Y) \cup C, Y}(1, d) \rightarrow 0. \tag{13}$$

From Formulæ (6) and (8), we have $h^0(\mathcal{O}_A(1, d)) = (d + 2)^2 = h^0(\mathcal{O}_F(1, d)) + 1$. Obviously we have that $B \cap Y$ is formed by $d + 1$ points, and up to the identification of Y with F_1 given in Formula (5), since the curve C corresponds to an element of type $h + f$ in F_1 , we can write

$$\mathcal{J}_{(B \cap Y) \cup C, Y}(1, d) \cong \mathcal{J}_{(B \cap Y) \cup C, F_1}(h + (d + 1)f) \cong \mathcal{J}_{B \cap Y, F_1}(df).$$

Since $A \in \mathcal{T}^*(d + 2)$ and every element of $|f|$ meets C (indeed $(h + f)f = 1$), the restriction to $B \cap Y$ of the ruling morphism $D \rightarrow \mathbb{P}^1$ associated with $|f|$

is injective. Thus, $h^1(Y, \mathcal{J}_{A \cap Y, Y}(1, d)) = 0$ and the exact sequence (13) gives $h^1(\mathcal{J}_B(1, d-1)) \geq h^1(\mathcal{J}_A(1, d)) \geq 2$, where the latter is greater than or equal to 2 because $\chi(\mathcal{J}_A(1, d)) = -1$ (see Formula (7)), and we assume that $h^0(\mathcal{J}_A(1, d)) \geq 1$. So we also have $h^0(\mathcal{J}_B(1, d-1)) > 0$.

Recall that $B \in \mathcal{T}^*(d+1)$ and thus, by the inductive assumption, B is not contained in any irreducible $E \in |\mathcal{O}_{\mathbb{F}}(1, d-1)|$. Therefore, thanks to Remarks 2.9 and 2.10, there must be an irreducible $M \in |\mathcal{O}_{\mathbb{F}}(1, 1)|$ containing at least 3 connected components of B , say $B' \subset M$ with B' of bidegree $(3, 3)$. Hence, by Lemma 3.1, there exists a curve L of bidegree $(1, 0)$ such that $\#(L \cap B') = 3$. So $A \notin \mathcal{T}^*(d+2)$, a contradiction. \square

Having proved that an irreducible bidegree $(1, d)$ -surface cannot contain $d+2$ (or more) non-collinear twistor fibers, we now prove that all the other cases can indeed occur. In particular, in the following result we prove a stronger version of Theorem 1.2 in the case of $n \leq d+1$.

Theorem 3.4. *Fix integers $d \geq 1$ and $0 \leq n \leq d+1$. Then, for any $A \in \mathcal{T}^*(n)$ there exists an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, d)|$ containing A . Moreover, the general $S \in |\mathcal{J}_A(1, d)|$ contains no other twistor fibers.*

Proof. We use induction on the integer d . If $d = 1$, the statement is true by Remark 3.2.

Now assume $d \geq 2$ and take an element $A \in \mathcal{T}^*(n)$. Since $\mathcal{T}^*(n)$ is Zariski dense in $\mathcal{C}(n)$, A has the bigraded Hilbert function of a general element of $\mathcal{C}(n)$. Thus, thanks to Corollary 2.20, we have that $h^1(\mathcal{J}_A(1, d)) = 0$ and, by Formula (7)

$$h^0(\mathcal{J}_A(1, d)) = (d+1)(d+3) - n(d+2) =: N_n + 1.$$

Fix a connected component C of A and set $B := A \setminus C$. If Y denotes the only $(0, 1)$ -surface containing C , Corollary 2.20 implies that $h^1(\mathcal{J}_B(1, d-1)) = 0$ and, again by Formula 7

$$h^0(\mathcal{J}_B(1, d-1)) = d(d+2) - (n-1)(d+1).$$

By the inductive assumption, we know that $|\mathcal{J}_B(1, d-1)| \neq \emptyset$ and a general $W \in |\mathcal{J}_B(1, d-1)|$ is irreducible. Thus $Y \cup W \in |\mathcal{J}_A(1, d)|$ and $Y \cup W$ has 2 irreducible components, one of which has bidegree $(1, d-1)$.

Let C_1, \dots, C_n be the connected components of A , $B_i := A \setminus C_i$ and Y_i be the unique element in $|\mathcal{J}_{C_i}(0, 1)|$. The set of all the reducible surfaces $W \cup Y_i \in |\mathcal{J}_A(1, d)|$, where $W \in |\mathcal{J}_{B_i}(1, d-1)|$, is the union of e projective spaces (one for each choice of C_i), each of them of codimension $h^0(\mathcal{J}_A(1, d)) - h^0(\mathcal{J}_B(1, d-1)) = d+2-n > 0$ in $|\mathcal{J}_A(1, d)| = \mathbb{P}^{N_n}$ (of codimension 1 if $n = d+1$). Therefore, they do not cover all $|\mathcal{J}_A(1, d)|$.

Now we want to exclude other possible splittings. In particular, we consider reducible surfaces of the form $W_1 \cup D_1$ with W_1 irreducible, D_1 possibly reducible of bidegree $(0, x)$ for some $x \geq 2$ and hence W_1 of bidegree $(1, d-x)$. Remark 2.10 shows that only irreducible components of D_1 of bidegree $(0, 1)$

may contain some component of A . We get that the surface is of the form $W_1 \cup D_2 \cup D_3$ with $W_1 \cup D_2$ of bidegree $(1, d - 1)$, but, as shown before, these kind of surfaces do not cover all $|J_A(1, d)|$, hence we have the thesis.

Now we prove that a general $S \in |J_A(1, d)|$ does not contain any other twistor fibers. We start by analyzing the case $d = 2$ and discussing the cases $n = 1, 2, 3$ separately.

Assume $n = 1$. Fix $C \in \mathcal{T}(1)$. Let $\mathcal{C}_C(1)$ denote the set of all $B \in \mathcal{C}(1)$ such that $B \cap C = \emptyset$. Note that $\mathcal{T}(1) \setminus \{C\} = \mathcal{C}_C(1) \cap \mathcal{T}(1)$. For any $B \in \mathcal{C}_C(1)$, we have $h^1(\mathcal{J}_{C \cup B}(1, 2)) = 0$ and hence $h^0(\mathcal{J}_{C \cup B}(1, 2)) = h^0(\mathcal{J}_C(1, 2)) - 4$. Let \mathcal{X}_C be the set of all smooth and irreducible surfaces of bidegree $(1, 2)$ containing C . It is a non-empty Zariski open subset of $|J_C(1, 2)|$. For any $B \in \mathcal{C}_C(1)$ the set of all $S \in \mathcal{X}_C$ containing B has complex codimension 4 and hence real codimension 8 as real manifolds. Since $\mathcal{T}(1)$ has real dimension 4, a general $S \in \mathcal{X}_C$ contains no other twistor fiber.

Now let $n = 2$. Fix $A \in \mathcal{T}(2)$ and let $\mathcal{C}_A(1)$ denote the set of all $B \in \mathcal{C}(1)$ such that $B \cap A = \emptyset$ and $B \cup A \in \mathcal{C}^*(3)$. Note that $h^1(\mathcal{J}_{A \cup B}(1, 2)) = 0$. So, as in the previous step, we get that a sufficiently general $S \in |J_A(1, 2)|$ does not contain an element of $\mathcal{T}_A(1)$. Assume S contains $B \in \mathcal{T}(1)$ such that there exists a curve of bidegree $(1, 0)$ that intersects every connected component of $A \cup B$. Note that L is uniquely determined by A . Let $\mathcal{C}(A, L)$ denote the set of all $B \in \mathcal{C}(1)$ such that $B \cap A = \emptyset$ and L meets B and set $\mathcal{T}(A, L) := \mathcal{C}(A, L) \cap \mathcal{T}(1)$. For any $o \in L \setminus (L \cap C)$, the set of all $B \in \mathcal{C}(A, L)$ containing o is a non-empty family of complex dimension 1, while there exists a unique twistor fiber containing o . The set $\mathcal{C}(A, L)$ is a complex manifold of dimension 2, while $h^1(\mathcal{J}_{A \cup B}(1, 2)) = 1$ and hence $h^0(\mathcal{J}_{A \cup B}(1, 2)) = h^0(\mathcal{J}_A(1, 2)) - 3$. Since $\dim \mathcal{C}(A, L) = 2$, a general $S \in |J_A(1, 2)|$ does not contain an element of $\mathcal{C}(A, L)$. Hence, there is no twistor fiber B such that $A \cup B \in \mathcal{T}^*(3)$.

Assume now that $n = 3$. First, consider a general $A \in \mathcal{T}^*(3)$. Then we have $h^1(\mathcal{J}_A(1, 2)) = 0$ and $h^0(\mathcal{J}_A(1, 2)) = 3$. For any $x \in \{0, 1, 2, 3\}$, let $\mathcal{C}(A, x)$ denote the set of all $C \in \mathcal{C}(1)$ such that $A \cap C = \emptyset$ and $h^0(\mathcal{J}_{A \cup C}(1, 2)) = x$ and set $\mathcal{T}(A, x) := \mathcal{C}(A, x) \cap \mathcal{T}(1)$. For a general $D \in \mathcal{C}(4)$ we have $h^0(\mathcal{J}_D(1, 1)) = 0$ (but $h^1(\mathcal{J}_D(1, 1)) = 1$). Fix $C \in \mathcal{C}(1)$ so that $C \cap A = \emptyset$, call Y the only element of $|J_C(0, 1)|$. Since $C \cap A = \emptyset$, no connected component of A is contained in Y , and $Y \cap A$ is formed by 3 points, all of them in $Y \setminus C$. We can easily handle the case $x = 0$, since any curve $C \in \mathcal{C}(A, 0)$ is not contained in any element of $|J_A(1, 2)|$. We have $\mathcal{O}_Y(1, 2)(-C) \cong \mathcal{O}_{F_1}(2f)$ and hence $C \in \mathcal{C}(A, 0)$ if no curve of bidegree $(1, 0)$ $L \in |f|$ intersects 2 of the components of A (since $h^1(\mathcal{J}_A(1, 1)) = 1$ the last statement is only an “if” and not an “if and only if”). A necessary condition for being $C \in \mathcal{C}(A, 2)$ is that L intersects all connected components of A , but this is excluded because $A \in \mathcal{T}^*(3)$. Since $A \in \mathcal{T}^*(3)$, there are exactly 3 curves L_1, L_2, L_3 of bidegree $(1, 0)$ intersecting 2 of the connected components of A . We claim that a general $S \in |J_A(1, 2)|$ does not contain $C \in \mathcal{C}(1)$ such that $C \cap A = \emptyset$. The family of smooth conics intersecting L_i has complex dimension 2, while the family of twistor fibers intersecting L_i has real dimension 2. As the

general $S \in |\mathcal{J}_A(1, 2)|$ has only finitely many conics, it has only a finite number of elements in $\mathcal{C}(A, 1)$ and, for the general, none of them is a twistor fiber.

We now proceed to analyze the case $d \geq 3$. Suppose the general surface of $|\mathcal{J}_A(1, d)|$ contains the twistor fiber $C \not\subseteq A$. So $A \cap C = \emptyset$. Set $A' := A \cup C$. Take $Y \in |\mathcal{O}_Y(0, 1)|$ containing C and consider the residual exact sequence

$$0 \rightarrow \mathcal{J}_A(1, d - 1) \rightarrow \mathcal{J}_{A'}(1, d) \rightarrow \mathcal{J}_{(Y \cap A) \cup C, Y}(1, d) \rightarrow 0. \tag{14}$$

We have $\mathcal{J}_{C, Y}(1, d) \cong \mathcal{O}_{F_1}(df)$.

Assume first $n \leq d$. If $A' \in \mathcal{T}^*(n + 1)$, then $h^1(\mathcal{J}'_A(1, d)) = 0$ and hence $h^0(\mathcal{J}'_A(1, d)) = h^0(\mathcal{J}_A(1, d)) - d - 2$. Since $\dim \mathcal{C}(1) = 4$, we have for $d \geq 3$ that the general $S \in |\mathcal{J}_A(1, d)|$ does not contain C , so $A \cup C \in \mathcal{T}^*(n + 1)$.

Now suppose $A' \notin \mathcal{T}^*(n + 1)$. Then there are connected components C' and C'' of A such that $C \cup C' \cup C'' \notin \mathcal{T}^*(3)$, i.e. C intersects the unique line L which meets C' and C'' . Since $\dim L = 1$, to exclude this case it is sufficient to prove that $h^0(\mathcal{J}_B(1, d)) \leq h^0(\mathcal{J}_A(1, d)) - 2$, i.e. $h^0(F_1, \mathcal{J}_{A \cap Y}(df)) \leq d$. But since $h^0(\mathcal{O}_{F_1}(df)) = d + 1$, then $\mathcal{O}_{F_1}(df)$ is globally generated and $A \cap Y \neq \emptyset$, therefore $h^0(F_1, \mathcal{J}_{A \cap Y}(df)) \leq d$.

Now let us assume $n = d + 1$ and that $A' \in \mathcal{C}^*(d + 2)$. By Lemma 2.21 we have $h^1(\mathcal{J}_{A'}(1, d)) \leq 1$ and hence $h^0(\mathcal{J}_{A'}(1, d)) \leq h^0(\mathcal{J}_A(1, d)) - d - 1$. Now suppose $A' \notin \mathcal{C}^*(d + 2)$. We need $h^0(F_1, \mathcal{J}_{A \cap Y}(df)) \leq d - 1$. Let $\pi : F_1 \rightarrow \mathbb{P}^1$ denote the ruling of F_1 . Since $\mathcal{O}_{\mathbb{P}^1}(d)$ is very ample, $h^0(F_1, \mathcal{J}_{A \cap Y}(df)) \leq d - 1$ if and only if $\#\pi(A \cap Y) \geq 2$, which is true because $\#(A \cap Y) = d + 1 \geq 3$ and (since $A \in \mathcal{C}^*(d + 1)$ no fiber F of π contains at least 3 points of A). The only remaining case is $d = 3$ and $n = 4$, which can be handled by simply adapting the previous argument for $d = 2$ and $n = 3$. \square

Having proved Theorem 1.2 in the case $n \leq d + 1$ for non collinear twistor fibers, we now focus on the case of $n = d + 2$. In this case we need some further preliminary results.

Lemma 3.5. *Fix $d > 0$, $n \leq d + 2$ and consider a general $A \in \mathcal{T}(n)$. Then $h^1(\mathcal{J}_A(1, d)) = 0$.*

Proof. Since $\mathcal{T}(n)$ is Zariski dense in $\mathcal{C}(n)$, it is sufficient to prove the statement for a general $A \in \mathcal{C}(n)$. Obviously, it is sufficient to prove the case $n = d + 2$. Take a connected component C of A and set $B := A \setminus C$ and $Y \in |\mathcal{J}_C(0, 1)|$. Consider the residual exact sequence of Y , as in Formula (13). Recall the correspondence $Y \cong F_1$ given in Formula (5) and let $\rho : Y \rightarrow \mathbb{P}^1$ denote its ruling. Since A is general $\#\rho(A \cap Y) = d + 1$ and hence $h^i(F_1, \mathcal{J}_{B \cap Y}(df)) = 0$, for $i \in \{0, 1\}$. Hence $h^1(\mathcal{J}_A(1, d)) = 0$. \square

From Theorem 1.1, we already know that an irreducible $(1, d)$ surface can contain a union of $d + 2$ twistor fibers only if it belongs to $\mathcal{T}(d + 2) \setminus \mathcal{T}^*(d + 2)$. Therefore, we introduce the following notation for sets of disjoint smooth conics that are collinear. Given a curve L of bidegree $(1, 0)$ and an integer $n > 0$, let $\mathcal{C}(n, L)$ denote the set of all $A \in \mathcal{C}(n)$ such that every connected component

of A meets L . The set $\mathcal{C}(n, L)$ is isomorphic (as real algebraic variety) to the set $\mathcal{S}(L, n)$ of all subsets of L with cardinality n and is therefore irreducible. An analogous definition and observation can be made for a curve R of bidegree $(0, 1)$, and of course, for the family \mathcal{T} instead of \mathcal{C} .

Lemma 3.6. *Fix integers $n \geq 3$ and $d \geq 1$ and take any $A \in \mathcal{T}(n, L)$. Then,*

$$h^1(\mathcal{J}_A(1, d)) \geq n - 2 + \max\{0, n - (d + 1)\}.$$

Proof. Let C_1, \dots, C_n be the connected components of A . Let $S_i \subset C_i$ be an arbitrary union of $d+2$ distinct points on each conic and $S := S_1 \cup \dots \cup S_n$. Since C_i is a smooth rational curve, the restriction map $H^0(\mathcal{O}_{C_i}(1, d)) \rightarrow H^0(\mathcal{O}_{S_i}(1, d))$ is bijective. So the restriction map

$$H^0(\mathcal{O}_A(1, d)) \rightarrow H^0(\mathcal{O}_S(1, d))$$

is bijective and $\chi(\mathcal{O}_A(1, d)) = \chi(\mathcal{O}_S(1, d))$. Thus $\chi(\mathcal{J}_A(1, d)) = \chi(\mathcal{J}_S(1, d))$.

Since $A \in \mathcal{T}(n, L)$, we know that there exists L of bidegree $(1, 0)$ which intersects every conic in A (and the $(0, 1)$ curve $j(L)$ does the same). So we can choose S such that n points are on L and n points are on $j(L)$. In other words we assume $(A \cap L) \cup (A \cap j(L)) \subseteq S$. Using Bézout and the fact that the bidegree is $(1, d)$, we get that $(n - 2) + \max\{0, n - (d + 1)\}$ of these points can be omitted without changing the set $|\mathcal{J}_S(1, d)|$, and hence $H^0(\mathcal{J}_S(1, d))$. It follows that $h^1(\mathcal{J}_A(1, d)) \geq n - 2 + \max\{0, n - (d + 1)\}$. \square

Remark 3.7. Thanks to the previous lemma, if $A \in \mathcal{T}(3, L)$, for some bidegree $(1, 0)$ curve L , then $h^1(\mathcal{J}_A(1, 1)) \geq 2$, and hence $h^0(\mathcal{J}_A(1, 1)) \geq 1$. However, since no surface of bidegree $(1, 0)$ or $(0, 1)$ contains an element of $\mathcal{T}(2)$, then every $S \in |\mathcal{J}_A(1, 1)|$ is irreducible. Hence, Proposition 2.17 gives that $|\mathcal{J}_A(1, 1)| = \{S\}$ and so, thanks to Formula (7), $h^1(\mathcal{J}_A(1, 1)) = 2$.

Finally, the following result completes the proof of Theorem 1.2, in the case of $d + 2$ twistor fibers.

Theorem 3.8. *Fix an integer $d \geq 2$ and take a general $A \in \mathcal{T}(d + 2, L)$. Then $h^0(\mathcal{J}_A(1, d)) \geq d$ and the general $S \in |\mathcal{J}_A(1, d)|$ is irreducible.*

Proof. Applying Lemma 3.6 with $n = d + 2$, we get $h^1(\mathcal{J}_A(1, d)) \geq d + 1$. Since $\chi(\mathcal{J}_A(1, d)) = (d + 1)(d + 3) - (d + 2)^2 = -1$, we get $h^0(\mathcal{J}_A(1, d)) \geq d$ and hence, $|\mathcal{J}_A(1, d)| \neq \emptyset$.

Now we prove that a general element in $|\mathcal{J}_A(1, d)|$ is irreducible. Take $S \in |\mathcal{J}_A(1, d)|$. Every surface of bidegree $(1, 0)$ or $(0, 1)$ contains at most one connected component of A . Therefore, S cannot be the union of a surface of bidegree $(1, 0)$ and d of bidegree $(0, 1)$. No irreducible and reduced surface of bidegree $(0, x)$, for $x \geq 2$, contains a twistor fiber. If $d = 2$, then $h^0(\mathcal{J}_A(1, 2)) \geq 2$. However, thanks to Remark 3.7, for any choice of $C \in A$ there is only one $M \in |\mathcal{J}_{(A \setminus C)}(1, 1)|$. Since we have considered all the possible reducible elements of $|\mathcal{J}_A(1, 2)|$, we get the thesis.

Now suppose $d > 2$. Since A has finitely many components, it suffices to prove that for any $x \in \{3, \dots, d - 1\}$, any union E of x connected components of A , and any connected component C of $A \setminus E$ we have

$$h^0(\mathcal{J}_E(1, x - 2)) < h^0(\mathcal{J}_{E \cup C}(1, x - 1)), \tag{15}$$

and then proceed as in the proof of Theorem 3.4.

Let Y be the only element of $|\mathcal{O}_{\mathbb{F}}(0, 1)|$ that contains C . The exact sequence in Formula (10) gives $h^0(\mathcal{J}_E(1, x - 2)) \leq h^0(\mathcal{J}_{E \cup C}(1, x - 1))$ and equality holds if and only if Y is in the base locus \mathcal{B} of $|\mathcal{J}_{E \cup C}(1, x - 1)|$. Call C_1, \dots, C_x the components of E and Y_i the only surface of bidegree $(0, 1)$ containing C_i . By Remark 2.13 the irreducible surfaces Y, Y_1, \dots, Y_x are all different from each other. For a general A we get that every integer $h^0(\mathcal{J}_E(1, x - 2))$ is the same for every union of x connected components of A . So if the inequality is false, then \mathcal{B} contains the surface $Y \cup Y_1 \cup \dots \cup Y_x$ of bidegree $(0, x + 1)$, which is a contradiction. \square

4. Surfaces of bidegree (1, 2) and (1, 3)

In this section we specialize our study to the case of surfaces of bidegree $(1, 2)$ and $(1, 3)$. In particular, we will prove Theorems 1.3, 1.4 and 1.5.

Recall, from Formula (7), that for each $A \in \mathcal{C}(n)$, we have $\chi(\mathcal{J}_A(1, 2)) = 15 - 4n$, and hence, if $n \leq 3$, we get $h^0(\mathcal{J}_A(1, 2)) > 0$. We also recall that a general $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ contains a finite number of smooth conics and, thanks to Theorem 2.19, for every $B \in \mathcal{C}(2)$ we have $h^1(\mathcal{J}_B(1, 1)) = 0$.

4.1. Surfaces of bidegree (1, 2) containing $0 \leq n \leq 4$ twistor fibers.

In this section, we show the existence of a smooth surface of bidegree $(1, 2)$ containing exactly $0, 1, 2, 3$ or 4 twistor fibers. In order to analyze the space $|\mathcal{J}_A(1, 2)|$ when A is in $\mathcal{C}(n)$ (or in $\mathcal{T}(n)$), for $0 \leq n \leq 4$, we need some preliminary results. Note that the extremal case, when $n = 4$, is treated differently.

We start by considering $(1, 2)$ -surfaces containing three disjoint smooth conics.

Proposition 4.1. *Take $A \in \mathcal{C}(3)$ such that $h^0(\mathcal{J}_A(1, 1)) > 0$. Then*

- (1) *there exists a curve L of bidegree $(1, 0)$ and a curve R of bidegree $(0, 1)$ such that $A \in \mathcal{C}(3, L)$ and $A \in \mathcal{C}(3, R)$*
- (2) *there is an irreducible element in $|\mathcal{J}_A(1, 2)|$;*
- (3) *$h^1(\mathcal{J}_A(1, 2)) = 1$;*
- (4) *the base locus \mathcal{B} of $|\mathcal{J}_A(1, 2)|$ is $A \cup L \cup R$, where L and R are the curves defined in (1).*

Proof. We start arguing as in Remark 3.7. Thanks to Remark 2.9, every surface of bidegree $(1, 0)$ or $(0, 1)$ does not contain an element of $\mathcal{C}(2)$, so as $A \in \mathcal{C}(3)$, every element in $|\mathcal{J}_A(1, 1)|$ is irreducible. Proposition 2.17 (with $a = b = c = d = 1$) gives that $h^0(\mathcal{J}_A(1, 1)) = 1$ and hence we set $|\mathcal{J}_A(1, 1)| = \{M\}$ and by Formula (7) we compute $h^1(\mathcal{J}_A(1, 1)) = 2$.

We now prove the first statement. Let C be a connected component of A , set $B := A \setminus C$ and denote by X the only element of $|J_C(1, 0)|$. Performing the same construction that leads to Formula (10), we have

$$0 \rightarrow \mathcal{J}_B(0, 1) \rightarrow \mathcal{J}_A(1, 1) \rightarrow \mathcal{J}_{A \cap X, X}(1, 1) \rightarrow 0.$$

Since $h^0(\mathcal{J}_B(0, 1)) = 0$ and $h^0(\mathcal{J}_A(1, 1)) = 1$, the previous residual exact sequence gives

$$h^0(\mathcal{J}_{A \cap X, X}(1, 1)) \geq 1. \tag{16}$$

Thanks to Formula (4), we have that $\mathcal{O}_X(1, 1) \simeq \mathcal{O}_{F_1}(h + 2f)$; moreover, recall from Remark 2.9 that C is identified with an element of $|\mathcal{O}_{F_1}(h + f)|$. Since $A \cap X$ is the union of C and the two points $B \cap X$, there exists a fiber $L \in |f|$ of the ruling of F_1 containing $B \cap X$. Since $f(h + f) = 1$, we have that L meets C . So L hits every connected component of A . If we take instead of X the only element of $|J_C(0, 1)|$, we get the existence of R .

We now prove (2). We will show that there is an irreducible element in $|J_A(1, 2)|$ by showing that the possible reducible cases do not cover the entire family. Remark 2.10 shows that A is not contained in a surface of bidegree $(1, 2)$ with an irreducible component of bidegree $(0, 2)$. By Remark 2.9, a bidegree $(0, 1)$ or $(0, 1)$ -surface does not contain any element of $\mathcal{C}(n)$, with $n \geq 2$. Thus, there are only finitely many elements of $|\mathcal{O}_{\mathbb{F}}(1, 2)|$ with at least 3 irreducible components.

Since $h^0(\mathcal{O}_{\mathbb{F}}(0, 1)) = 3$, the set of all reducible elements of $|\mathcal{O}_{\mathbb{F}}(1, 2)|$ with an irreducible component of bidegree $(1, 1)$ containing A is isomorphic to \mathbb{P}^2 . Thus, in order to prove the existence of an irreducible element in $|J_A(1, 2)|$, it is sufficient to show that $h^0(\mathcal{J}_A(1, 2)) \geq 4$. But, using the exact sequence (9), this is equivalent to proving that $h^1(\mathcal{J}_A(1, 2)) \geq 1$, and the last inequality is true, by Theorem 2.19 because $\#(L \cap A) = 3$.

To prove (3), i.e. $h^1(\mathcal{J}_A(1, 2)) = 1$, it is sufficient to prove that we have $h^1(\mathcal{J}_A(1, 2)) \leq 1$. As before, take a connected component C of A and set $B = A \setminus C$. Let Y be the only element of $|J_C(0, 1)|$. In the identification (5) of Y with F_1 we have

$$\mathcal{J}_{(B \cap Y) \cup C, Y}(1, 2) \cong \mathcal{J}_{(B \cap Y) \cup C, F_1}(h + 3f) \cong \mathcal{O}_{(B \cap Y), F_1}(2f).$$

Since $\#(B \cap Y) = 2$ and $\mathcal{O}_{F_1}(2f)$ is globally generated, $h^1(F_1, \mathcal{J}_{B \cap Y, F_1}(2f)) \leq 1$. We have $\text{Res}_Y(A) = B$. Thanks to Theorem 2.19, we have $h^1(\mathcal{J}_B(1, 1)) = 0$, and the residual exact sequence of Y ,

$$0 \rightarrow \mathcal{J}_B(1, 1) \rightarrow \mathcal{J}_A(1, 2) \rightarrow \mathcal{J}_{(B \cap Y) \cup C, Y}(1, 2) \rightarrow 0,$$

gives $h^1(\mathcal{J}_A(1, 2)) \leq 1$.

Finally, we discuss the base locus of $|J_A(1, 2)|$ in order to prove (4). First of all, for any surface $S \in |J_A(1, 2)|$, we clearly have $A \subset S$. Moreover, since $\#(L \cap A) = 3$ and $\#(R \cap A) = 3$, then by Bézout, both curves are contained in S : in fact, thanks to Remark 2.3 and Formula (3), the general intersection between a curve of bidegree $(1, 0)$ and S consists of one point while the intersection of

a curve of bidegree $(0, 1)$ and S consists of two points (see also Remark 2.16). Therefore, $L \cup R \subset S$ and $A \cup L \cup R \subset \mathcal{B}$.

We now prove that $\mathcal{B} \subset A \cup L \cup R$. Fix $p \in \mathcal{B} \setminus (A \cup L \cup R)$. Take a connected component $C_i, i = 1, 2, 3$, of A and set $B_i := A \setminus C_i$. Let Y_i be the only element of $|\mathcal{O}_{\mathbb{F}}(0, 1)|$ containing C_i . By Proposition 2.22 $B_i \cup L \cup R$ is the base locus of $|\mathcal{J}_{B_i}(1, 1)|$. So there exists $S_i \in |\mathcal{J}_{B_i}(1, 1)|$ such that $p \notin S_i$. If $p \notin Y_i$, then $p \notin \mathcal{B}$. Since $S_1 \cap S_2 \cap S_3 = L \cup R$, we can take $i \in \{1, 2, 3\}$ such that $p \notin Y_i$. So $\mathcal{B} = A \cup L \cup R$. \square

The following remark shows that if $A \in \mathcal{C}(3)$ satisfies condition (1) of Theorem 4.1, then the existence of a $(1, 1)$ -surface containing A is guaranteed. In particular there exists a $(1, 1)$ -surface containing arbitrary triplets of collinear twistor fibers.

Remark 4.2. Take $A \in \mathcal{C}(3)$ and assume the existence of curves L of bidegree $(1, 0)$ and R of bidegree $(0, 1)$ intersecting every connected component of A . By adapting the proof of Lemma 3.6, since $\#(L \cap A) = 3$ and $\#(R \cap A) = 3$, we have that $h^1(\mathcal{J}_A(1, 1)) \geq 2$. Thus $h^0(\mathcal{J}_A(1, 1)) \geq 1$ and A satisfies the assumptions of Proposition 4.1.

We can be even more specific and say that if $A \in \mathcal{C}(3)$ (with no assumption about L or R), then $h^0(\mathcal{J}_A(1, 1)) \leq 1$ and if $|\mathcal{J}_A(1, 1)| \neq \emptyset$, then the only element of $|\mathcal{J}_A(1, 1)|$ is irreducible. This is true because, thanks to Remark 2.9, every reducible element of $|\mathcal{O}_{\mathbb{F}}(1, 1)|$ contains at most 2 disjoint smooth conics.

Note that if $A \in \mathcal{T}(3)$ and L exists, then we can take $R := j(L)$. So if $A \in \mathcal{T}(3)$ to get $h^0(\mathcal{J}_A(1, 1)) > 0$, it is sufficient to assume that $A \notin \mathcal{T}^*(3)$.

The following lemma is a sort of inverse of the previous remark.

Lemma 4.3. *Take $A \in \mathcal{C}^*(3)$. Then $h^0(\mathcal{J}_A(1, 1)) = 0$ and $h^1(\mathcal{J}_A(1, 1)) = 1$.*

Proof. If $A \in \mathcal{C}(3)$, thanks to Formula (7), $\chi(\mathcal{J}_A(1, 1)) = -1$. So we have $h^0(\mathcal{J}_A(1, 1)) = 0$ if and only if $h^1(\mathcal{J}_A(1, 1)) = 1$. We assume that $h^0(\mathcal{J}_A(1, 1)) \neq 0$ and will prove that $A \notin \mathcal{C}^*(3)$. Let $B \subset A$ be the union of 2 connected components of A and set $C := A \setminus B$. Let L and R be the curves defined in Remark 2.5 for $B \in \mathcal{C}(2)$. Take any element $D \in |\mathcal{J}_B(1, 1)|$. Since $\#(B \cap (L \cup R)) = 2, B \subset D$, and D has bidegree $(1, 1)$, then the Bézout theorem implies $L \cup R \subset D$.

By Theorem 2.19 and Proposition 2.22, we have that

$$h^1(\mathcal{J}_B(1, 1)) = 0, \quad h^0(\mathcal{J}_B(1, 1)) = 2,$$

and the general element M in $|\mathcal{J}_B(1, 1)|$ is irreducible. Since $h^0(\mathcal{J}_B(1, 1)) = 2$ and M is general, $C \not\subset M$. Consider the following residual exact sequence:

$$0 \rightarrow \mathcal{J}_C \rightarrow \mathcal{J}_A(1, 1) \rightarrow \mathcal{J}_{B \cup (M \cap C), M}(1, 1) \rightarrow 0 \tag{17}$$

Since $M \in |\mathcal{J}_B(1, 1)|$ and $h^1(\mathcal{O}_{\mathbb{F}}) = 0$, the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{F}} \rightarrow \mathcal{J}_B(1, 1) \rightarrow \mathcal{J}_{B, M}(1, 1) \rightarrow 0$$

gives $h^0(M, \mathcal{J}_{B, M}(1, 1)) = 1$. Moreover, $h^0(\mathcal{J}_C) = 0$, and so the sequence (17) and the assumption $h^0(\mathcal{J}_A(1, 1)) \geq 1$ imply $h^0(M, \mathcal{J}_{B \cup (M \cap C), M}(1, 1)) \geq 1$. By

Proposition 2.22, the curve $A \cup L \cup R$ is the base locus of $|J_B(1, 1)|$ and thus the base locus of $H^0(M, \mathcal{J}_{B,M}(1, 1))$ is the curve $B \cup L \cup R$. Since $B \cap C = \emptyset$, the degree 2 scheme $C \cap M$ is contained in $L \cup R$. To get $A \notin \mathcal{C}^*(3)$, we must to prove that $C \cap L \neq \emptyset$. It is sufficient to observe that $\deg(C \cap T) \leq 1$ for any curve T of bidegree $(0, 1)$. This is true because of Remark 2.3 and the fact that C is the intersection of a surface of bidegree $(1, 0)$ and a surface of bidegree $(0, 1)$. \square

We now discuss the case of $A \in \mathcal{C}(2)$ contained in a smooth surface of bidegree $(1, 1)$. In this case we will also prove smoothness for the general element in $|J_A(1, 2)|$.

Proposition 4.4. *Take any $A \in \mathcal{C}(2)$ that is contained in a smooth element of $|\mathcal{O}_{\mathbb{F}}(1, 1)|$. Then we have*

- (1) $h^1(J_A(1, 2)) = 0$ and $h^0(J_A(1, 2)) = 7$;
- (2) the set $A \cup L$ is contained in the base locus \mathcal{B} of $|J_A(1, 2)|$, where L is the bidegree $(1, 0)$ curve described in Remark 2.5;
- (3) a general $S \in |J_A(1, 2)|$ is smooth.

Proof. To prove part (1) it is sufficient to apply Corollary 2.20, which gives $h^1(J_A(1, 2)) = 0$ and Formula (7), which entails $h^0(J_A(1, 2)) = 7$.

We now pass to point (2). Take L and R as in Remark 2.5. Since $\mathcal{O}_{\mathbb{F}}(0, 1)$ is globally generated, $\mathcal{B} \subseteq A \cup L \cup R$; moreover, thanks to Remark 2.16, we also have $A \cup L \subseteq \mathcal{B}$.

We are left to prove (3). By Bertini’s theorem $\text{Sing}(S) \subseteq A \cup L \cup R$ for a general $S \in |J_A(1, 2)|$. Fix a smooth $M \in |J_A(1, 1)|$. Take a general $Y' \in |\mathcal{O}_{\mathbb{F}}(0, 1)|$. Since Y' is general $L \cap Y' = \emptyset$ (and hence it is not singular at any $p \in L$). Thus, up to small deformation, we can say that S (which is general) is smooth in a neighborhood of L . We are left to exclude the case $\text{Sing}(S) \subseteq A \cup R$. Fix $p \in A \cup R$ and let $2p$ be the 0-dimensional scheme of \mathbb{F} defined by the ideal $\mathcal{J}_{p, \mathbb{F}}^2$. S is singular at p if and only if $S \in |J_{2p \cup A}(1, 2)|$. To complete our proof we need to prove that

$$h^0(J_{2p \cup A}(1, 2)) = h^0(J_A(1, 2)) - 2,$$

for all $p \in (A \cup R) \setminus A \cap R$ and that, for $p \in A \cap R$, $h^0(J_{2p \cup A}(1, 2)) < h^0(J_A(1, 2))$. These two statements give the thesis because $(A \cup R) \setminus A \cap R$ and $A \cap R$ are 1-dimensional and 0-dimensional, respectively, and we are saying that the set of bidegree $(1, 2)$ -surfaces containing A and a singular point has codimension 2 in the first case and positive codimension in the second one.

Let us start by taking $p \in (A \cup R) \setminus A \cap R$. Since p is a smooth point of $A \cup R$, $\deg(2p \cap (A \cup R)) = 2$. Consider the exact sequence

$$0 \rightarrow \mathcal{J}_{(A \cup R) \cup 2p}(1, 2) \rightarrow \mathcal{J}_{A \cup R}(1, 2) \rightarrow \mathcal{J}_{A \cup R} \otimes \mathcal{O}_{2p}(1, 2) \rightarrow 0. \tag{18}$$

Since $\deg(2p) = 4$ and A is smooth, we have $h^0(\mathcal{J}_{A \cup R} \otimes \mathcal{O}_{2p}(1, 2)) = 2$ if $p \in A \cup R$ and $h^0(\mathcal{J}_A \otimes \mathcal{O}_{2p}(1, 2)) = 4$ if $p \in R$. Hence it is sufficient to prove that

$$h^1(\mathcal{J}_{(A \cup R) \cup 2p}(1, 2)) = 0.$$

First of all, assume that $p \in A \setminus R$. Let C be the connected component of A containing p . Set the following notation $E := A \setminus C$. Since R is in the base locus of $\mathcal{J}_A(1, 1)$ we have that $h^0(\mathcal{J}_A(1, 1)) = h^0(\mathcal{J}_{A \cup R}(1, 1))$ (see [3, proof of Theorem 1.1]). Moreover, thanks to part (1) and to [3, Remark 4.3], we have $h^0(\mathcal{J}_{A \cup R}(1, 1)) = h^0(\mathcal{J}_A(1, 1)) = h^0(\mathcal{J}_E(1, 1)) - 3$. Thus p is not in the base locus of $|\mathcal{J}_E(1, 1)|$. Fix $M \in |\mathcal{J}_E(1, 1)|$ such that $p \notin S$. Let Y be the surface of $|\mathcal{O}_{\mathbb{F}}(0, 1)|$ containing C and consider the residual exact sequence with respect to Y :

$$0 \rightarrow \mathcal{J}_{E \cup p}(1, 1) \rightarrow \mathcal{J}_{A \cup 2p}(1, 2) \rightarrow \mathcal{J}_{(E \cap Y) \cup C \cup (2p \cap Y), Y}(1, 2) \rightarrow 0. \tag{19}$$

We now prove that

$$h^1(\mathcal{J}_{E \cup p}(1, 1)) = 0. \tag{20}$$

Recall that $A = E \cup C$ and $p \in C$, hence we have the exact sequence

$$0 \rightarrow \mathcal{J}_A(1, 1) \rightarrow \mathcal{J}_{E \cup p}(1, 1) \rightarrow \mathcal{J}_{p, C}(2) \rightarrow 0.$$

Thanks to Theorem 2.19 we have that $h^1(\mathcal{J}_A(1, 1)) = 0$; on the other hand, since C is a smooth rational curve, we have $h^1(\mathcal{J}_{p, C}(2)) = h^1(\mathcal{O}_C(1)) = 0$, and this proves (20).

To conclude, it is now sufficient to prove that

$$h^1(\mathcal{J}_{(E \cap Y) \cup C \cup (2p \cap Y), Y}(1, 2)) = 0. \tag{21}$$

Note that $\mathcal{J}_{C, Y}(1, 2) \cong \mathcal{O}_{F_1}(2f) \cong \mathcal{O}_Y(0, 1)$, hence, by [3, Remark 2.11] we know that $\mathcal{J}_{C, Y}(1, 2)$ is very ample. Therefore we get $h^1(\mathcal{J}_{C \cup (2p \cap Y), Y}(1, 2)) = 0$. Since $E \cap Y$ consists of one point, we conclude that (21) holds.

Thus the exact sequence (19) gives $h^1(\mathcal{J}_{A \cup 2p, \mathbb{F}}(1, 2)) = 0$, completing the proof in the case $p \in A \setminus R$.

Fix $p \in R \setminus (A \cap R)$ and recall that we need to prove that $h^0(\mathcal{J}_{A \cup 2p}(1, 2)) = 5$. Fix a general $Y' \in |\mathcal{J}_p(0, 1)|$. Since Y' is general, $R \not\subseteq Y'$ (and also $L \not\subseteq Y'$). Since Y' is smooth, $Y' \cap 2p = (2p, Y')$ is a scheme of degree 3 and $\text{Res}_{Y'}(2p) = \{p\}$. As $p \in R$, we have that $h^0(\mathcal{J}_{A \cup \{p\}}(1, 1)) = h^0(\mathcal{J}_A(1, 1)) = 2$. Thus, by the residual exact sequence of Y' it is sufficient to prove that

$$h^0(Y', \mathcal{J}_{(A \cap Y') \cup (2p, Y')}(1, 2)) \leq 3.$$

Since $\mathcal{O}_{Y'}(1, 2)$ is very ample, we have $h^0(Y', \mathcal{J}_{(2p, Y')}(1, 2)) = h^0(Y', \mathcal{O}_{Y'}(1, 2)) - 3 = 4$. So it is enough to prove that $A \cap Y'$ is not contained in the base locus, \mathcal{B}' , of $|\mathcal{O}_{(2p, Y')}(1, 2)|$. In the identification between Y' and F_1 we have $\mathcal{O}_{Y'}(1, 2) \cong \mathcal{O}_{F_1}(h + 3f)$. Let N be the only element of $|f|$ that contains p . We have $N \cong \mathbb{P}^1$ and $h^1(N, \mathcal{J}_{2p \cap N}(1, 2)) = 0$, but $N \subseteq \mathcal{B}'$. Since $\mathcal{O}_{F_1}(h + 2f)$ is very ample, $\mathcal{J}_p(h + 2f)$ has only p in its base locus. Thus $\mathcal{B}' = N$ and so, since $R \not\subseteq Y'$ (and also $L \not\subseteq Y'$), \mathcal{B}' cannot contain both points of $A \cap Y'$.

The last case is $p \in A \cap R$. To prove our claim, i.e. that $h^0(\mathcal{J}_{2p \cup A}(1, 2)) < h^0(\mathcal{J}_A(1, 2))$, it suffices to use (18). Hence, a general S is smooth. \square

The following result is analogous to Proposition 4.1, if we choose the conics to be twistor fibers.

Proposition 4.5. *Take $A \in \mathcal{T}(3)$ such that $h^0(\mathcal{J}_A(1, 1)) = 0$. Then we have the following:*

- (1) $h^1(\mathcal{J}_A(1, 2)) = 0$ and hence $h^0(\mathcal{J}_A(1, 2)) = 3$;
- (2) *there is an irreducible $S \in |\mathcal{J}_A(1, 2)|$;*
- (3) *the base locus of $|\mathcal{J}_A(1, 2)|$ is contained in the union of A and 3 different curves of bidegree $(1, 0)$;*
- (4) *for a sufficiently general A (contained in a dense euclidean open subset of $\mathcal{T}(3)$), we can take a smooth $S \in |\mathcal{J}_A(1, 2)|$.*

In the previous statement we assume $A \in \mathcal{T}(3)$ such that $h^0(\mathcal{J}_A(1, 1)) = 0$. Note that, thanks to Lemma 3.1, this implies that the conics in A do not belong to any infinite family of twistor fibers contained in a smooth j -invariant surface of bidegree $(1, 1)$.

Proof. We start with the proof of (1). Fix a connected component C of A and call D the only element of $|\mathcal{J}_C(0, 1)|$. Set $B := A \setminus C$. To get $h^1(\mathcal{J}_A(1, 2)) = 0$, mimicking the proof of Proposition 4.1 it is sufficient to prove $h^1(F_1, \mathcal{J}_{B \cap D}(2f)) = 0$. Suppose $h^1(F_1, \mathcal{J}_{B \cap D}(2f)) > 0$, i.e. assume the existence of $T \in |\mathcal{O}_{F_1}(f)|$ containing the 2 points $B \cap D$. Since $C \in |\mathcal{O}_{F_1}(h+f)|$, $C \cap T \neq \emptyset$. So T hits every connected component of A . Remark 4.2 gives $h^0(\mathcal{J}_A(1, 1)) > 0$, a contradiction.

To prove (2), it is sufficient to show that the reducible cases do not cover the whole $|\mathcal{J}_A(1, 2)|$. In fact, reasoning as in the proof of Proposition 4.1, the only possible splittings are of the form $(1, 0) + (0, 1) + (0, 1)$, which are in a finite number, or $(1, 1) + (0, 1)$, where the bidegree $(1, 1)$ component contains 2 connected components of A and the remaining bidegree $(0, 1)$ part is uniquely determined. Now, $h^0(\mathcal{J}_B(1, 1)) = 2$, so, the set of all reducible elements of $|\mathcal{J}_A(1, 2)|$ with an irreducible component of bidegree $(1, 1)$ does not cover $|\mathcal{J}_A(1, 2)|$.

We now prove (3) and (4). Since $h^0(\mathcal{J}_A(1, 1)) = 0$ and A is j -invariant, neither $\pi_1(A)$ nor $\pi_2(A)$ has a triple point (both have 3 double points). Set $L_1 \cup L_2 \cup L_3 := \pi_2^{-1}(\text{Sing}(\pi_2(A)))$ and $R_1 \cup R_2 \cup R_3 := \pi_1^{-1}(\text{Sing}(\pi_1(A)))$. Since $\#(L_i \cap A) = \#(R_i \cap A) = 2$, $L_1 \cup L_2 \cup L_3$ are in the base locus of $|\mathcal{J}_A(1, 2)|$ and each L_i and each R_i meets exactly 2 connected components of A .

To prove the existence of a smooth element, it is sufficient to reason as in the proof of Proposition 4.4 case (3). □

We are now ready to prove the first part of Theorem 1.3.

Theorem 4.6. *Fix $n \in \{0, 1, 2, 3\}$. There is a smooth $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ which contains exactly n twistor fibers.*

Proof. A general $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ contains only finitely many smooth conics. Since the set of all twistor fibers has real codimension 4 in the space of all smooth conics, a general $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ contains no twistor fiber.

Now we prove the case $n = 1$. Fix a twistor fiber C and take a general $S \in |\mathcal{J}_C(1, 2)|$. Assume that S contains another twistor fiber, E . We have $h^1(\mathcal{J}_C(1, 2)) = h^1(\mathcal{J}_{C \cup E}(1, 2)) = 0$ (Theorem 2.19 and Remark 2.14). Thus $|\mathcal{J}_{C \cup E}(1, 2)|$ is a 4-codimensional complex projective subspace of $|\mathcal{J}_C(1, 2)|$ (this

is explained by the equality $h^0(\mathcal{J}_{C \cup E}(1, 2)) = h^0(\mathcal{J}_C(1, 2)) - 4$ contained in [3, proof of Theorem 1.1]). However $\mathcal{T}(1)$ is a real 4-dimensional space. So a general $S \in |\mathcal{J}_C(1, 2)|$ does not contain any other twistor fiber.

Note that C is the base locus of $|\mathcal{J}_C(1, 2)|$. By Bertini’s theorem a general $S \in |\mathcal{J}_C(1, 2)|$ is smooth outside C . Fix $p \in C$ and let $2p$ be the closed subscheme of \mathbb{F} with $(\mathcal{J}_p)^2$ as its ideal sheaf. Recall that $2p \subset S$ if and only if $p \in \text{Sing}(S)$. Since $\dim C = 1$, to get that S is smooth, it is sufficient to prove that $h^0(\mathcal{J}_{2p \cup C}(1, 2)) \leq h^0(\mathcal{J}_C(1, 2)) - 2 = 9$. This follows from the proof of Proposition 4.4 case (3).

The case $n = 2$ is true by Proposition 4.4 with $\mathcal{T}(2)$ instead of $\mathcal{C}(2)$.

The case $n = 3$ is true by Proposition 4.5. □

In the remainder of this section, we will construct a smooth $(1, 2)$ -surface containing 4 twistor fibers. The following lemma, in the case $d = 2$, says that if an irreducible $(1, 2)$ -surface contains 4 disjoint smooth conics, then these conics are not general, because three of them must be collinear.

Lemma 4.7. *Let $d \geq 2$ and $A \in \mathcal{C}(d+2)$. If there is an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, d)|$, then $A \notin \mathcal{C}^*(d + 2)$.*

Proof. We prove the lemma by induction on d . We start with the case $d = 2$. Suppose that $A \in \mathcal{C}^*(4)$, i.e. there is no union B of 3 of the connected components of A such that $\#(L \cap B) = 3$ for some curve L of bidegree $(1, 0)$. Fix a connected component C of A and set $B := A \setminus C$. Call Y the only element of $|\mathcal{J}_C(0, 1)|$. Remark 2.9 gives $\text{Res}_Y(A) = B$. By assumption and Lemma 4.3, $h^0(\mathcal{J}_B(1, 1)) = 0$. Since $h^0(\mathcal{J}_A(1, 2)) \neq 0$, the residual exact sequence

$$0 \rightarrow \mathcal{J}_B(1, 1) \rightarrow \mathcal{J}_A(1, 2) \rightarrow \mathcal{J}_{A \cap Y, Y}(1, 2) \rightarrow 0,$$

gives $h^0(Y, \mathcal{J}_{A \cap Y, Y}(1, 2)) > 0$ (otherwise $|\mathcal{J}_A(1, 2)| = \emptyset$). The scheme $A \cap Y$ is the union of C and the 3 points $B \cap Y$. In the identification of Y with F_1 the line bundle $\mathcal{O}_Y(1, 2)$ goes to the line bundle $\mathcal{O}_{F_1}(h + 3f)$ and C goes to an element of $|h + f|$. Thus $h^0(F_1, \mathcal{J}_{B \cap Y, F_1}(2f)) > 0$. Thus at least 2 of the 3 points $B \cap Y$ are in the same fiber \hat{L} of the ruling $|f|$ of F_1 . Since $\hat{L} \cap C \neq \emptyset$, \hat{L} is a curve of bidegree $(1, 0)$ that meets at least 3 connected components of A . Let B' be the union of 3 components of A that intersect \hat{L} . The curves B' and \hat{L} give a contradiction.

Now assume that the result is true for $d + 1$. Note that as a by-product of the previous part, if $B \in \mathcal{C}^*(d + 1)$, then $h^0(\mathcal{J}_B(1, d - 1)) = 0$.

Assume $A \in \mathcal{C}^*(d + 2)$ and that there is an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, d)|$. Fix a connected component C of A and set $B := A \setminus C$. Take a surface Y of bidegree $(0, 1)$ containing C . By means of the sequence in Formula (10), we either have $h^0(\mathcal{J}_B(1, d - 1)) > 0$ or $h^0(Y, \mathcal{J}_{A \cap Y, Y}(1, d)) > 0$. Since $A \in \mathcal{C}^*(d + 2)$, $B \in \mathcal{C}^*(d + 1)$ and hence, thanks to the inductive assumption, we have $h^0(\mathcal{J}_B(1, d - 1)) = 0$. The scheme $A \cap Y$ is the union of C and the scheme $B \cap Y$ with $A \cap B \cap Y = \emptyset$. Up to the identification of Y and F_1 we have $\mathcal{O}_Y(1, d)(-C) \cong \mathcal{O}_{F_1}(df)$. Since Y has bidegree $(0, 1)$, every connected component of B is either contained in Y or it intersects transversely Y at a unique point. By Remark 2.9,

the set $B \cap Y$ is formed by $d + 1$ points. Thus $J_{A \cap Y, Y}(1, d) \cong J_{B \cap Y}(df)$. We saw that $h^0(Y, J_{A \cap Y, Y}(1, d)) > 0$, and this is true if and only if there are $u_1, \dots, u_{d+1} \in B \cap Y$ and $F \in |f|$ such that $u_i \neq u_j$, for $i \neq j$ and $\{u_1, \dots, u_{d+1}\} \subset F$. The set $F \cap C$ is a unique point, o , and $o \notin \{u_1, \dots, u_{d+1}\}$, because $B \cap C = \emptyset$. The curve F has bidegree $(0, 1)$ and therefore $A \notin \mathcal{C}^*(d + 2)$, a contradiction. \square

Thanks to the previous result, if an irreducible $(1, 2)$ -surface contains 4 disjoint smooth conics, then they are in special position. We now show that if these 4 conics are twistor fibers, then their position is very special. We start by introducing the following notation. For $n \geq 4$, we denote by $\mathcal{C}(n)^-$ the set of the elements $A \in \mathcal{C}(n)$ for which there exists a bidegree $(1, 0)$ curve L such that $A \in \mathcal{C}(n, L)$. The set $\mathcal{C}(n)^-$ parametrizes the families of n collinear disjoint smooth conics. For $n \geq 4$ we also write $\mathcal{T}(n)^- := \mathcal{T}(n) \cap \mathcal{C}(n)^-$. The families $\mathcal{C}(n)^-$ and $\mathcal{T}(n)^-$ are Zariski closed in $\mathcal{C}(n)$ and $\mathcal{T}(n)$, respectively.

The following lemma shows that if an irreducible $(1, 2)$ -surface contains 4 twistor fibers, then they are all collinear.

Lemma 4.8. *Take an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ containing $A \in \mathcal{T}(4)$. Then $A \in \mathcal{T}(4)^-$.*

Proof. Suppose there exists an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ containing $A \in \mathcal{T}(4)$. By Lemma 4.7 there exists a union B of 3 of the connected components of A such that $B \in \mathcal{T}(3) \setminus \mathcal{T}^*(3)$, i.e., there exists a bidegree $(1, 0)$ curve L , such that $B \in \mathcal{T}(3, L)$ and thus, thanks to Remark 3.7 $h^0(J_B(1, 1)) > 0$. However, the same remark tells us that $h^0(J_B(1, 1)) = 1$, $h^1(J_B(1, 1)) = 2$, and that the only element M of $|J_B(1, 1)|$ is irreducible.

As usual, set $C := A \setminus B$. As in Remark 4.2 since L of bidegree $(1, 0)$ meets every connected component of B , then $R := j(L)$, of bidegree $(0, 1)$, do the same.

Thanks to Remark 2.16 we get $B \cup L \cup R \subset M$. Since S and M are irreducible, thanks to Lemma 2.2, the one-dimensional scheme $S \cap M$ has bidegree $(5, 4)$. Since $B \cup L \cup R$ has bidegree $(4, 4)$, then $C \not\subset M$. Let Y be only element of $|\mathcal{J}_C(0, 1)|$. Since $B \subset M \cup Y$, then $M \cup Y \in |\mathcal{J}_A(1, 2)|$. Moreover, since S is irreducible, then $S \neq M \cup Y$, and hence $h^0(\mathcal{J}_A(1, 2)) \geq 2$, i.e. $h^1(\mathcal{J}_A(1, 2)) \geq 3$. Since $h^1(J_B(1, 1)) = 2$, the residual exact sequence

$$0 \rightarrow J_B(1, 1) \rightarrow J_A(1, 2) \rightarrow J_{A \cap Y, Y}(1, 2) \rightarrow 0,$$

gives $h^1(Y, J_{A \cap Y, Y}(1, 2)) > 0$. As in the proof of Lemma 4.7 we obtain the following inequality $h^1(F_1, J_{B \cap Y}(2f)) > 0$, i.e. there exists a curve $\hat{L} \in |f|$ of bidegree $(1, 0)$ which intersects at least 2 of the connected components of B .

Call B' the union of 2 of the connected components of B intersecting \hat{L} . Since $\hat{L} \cap C \neq \emptyset$ and $B' \cup C$ is j -invariant, every connected component of $B' \cup C$ meets $j(\hat{L})$. Remark 4.2, Proposition 4.1 and Bézout imply the existence of an irreducible surface M' of bidegree $(1, 1)$ containing $B' \cup C \cup \hat{L} \cup j(\hat{L})$. Since $B' \subset M'$, \hat{L} and $j(\hat{L})$ contain at least 2 points of M' , then $B' \cup \hat{L} \cup j(\hat{L}) \subset M'$. But by Remark 2.5 there is a unique curve of bidegree $(1, 0)$ intersecting two

different smooth conics, so $\hat{L} = L$ and both L and $j(L)$ intersect every connected component of B . Thus L intersects each connected component of A , i.e. $A \in \mathcal{C}(4)^-$. \square

As a byproduct of the proof of the previous result, we get the following lemma, which says that there are infinitely many irreducible $(1, 2)$ -surfaces containing 4 collinear twistor fibers.

Lemma 4.9. *Take $A \in \mathcal{T}(4)^-$ and assume that A is not contained in a surface of bidegree $(1, 1)$. Then $\dim |J_A(1, 2)| = 1$ and $|J_A(1, 2)|$ contains exactly 4 reducible elements of $|\mathcal{O}_{\mathbb{F}}(1, 2)|$.*

Proof. Let L be the curve of bidegree $(1, 0)$ that intersects every connected component of A . Since every connected component of A is j -invariant, $j(L)$ intersects every connected component of A . In the proof of Lemma 4.8 we showed that $h^0(J_A(1, 2)) \geq 2$. From the lines of that proof, it can be deduced that only 4 elements in $|J_A(1, 2)|$ are reducible and they are all obtained by fixing a connected component C of A and taking the union of the unique surface M_C of bidegree $(1, 1)$ containing $A \setminus C$ and the unique surface Y_C of bidegree $(0, 1)$ containing C . To complete the proof, it is sufficient to show that $h^0(J_A(1, 2)) \leq 2$. Take a connected component C of A and consider the residual exact sequence

$$0 \rightarrow J_C(0, 1) \rightarrow J_A(1, 2) \rightarrow J_{M_C \cap A, M_C}(1, 2) \rightarrow 0. \tag{22}$$

We have $h^0(J_C(0, 1)) = 1$, because the intersection of 2 different elements of $|\mathcal{O}_Y(0, 1)|$ is a curve of bidegree $(1, 0)$. Thus by (22) to conclude the proof it is sufficient to prove that the image \mathcal{V} of $H^0(J_A(1, 2))$ in $H^0(M_C, J_{M_C \cap A, M_C}(1, 2))$ has dimension at most 1. Bezout gives that $A \cup L \cup j(L)$ is contained in the base locus of $|J_{B, M_C}(1, 2)|$. Every $D \in |\mathcal{V}|$ has bidegree $(5, 4)$ as a curve of \mathbb{F} and thus a general $D \in |\mathcal{V}|$ is the union (counting multiplicities as divisors of the smooth surface M_C) of $A \cup L \cup j(L)$ and a curve E of bidegree $(1, 0)$ as a curve of \mathbb{F} . Recall that M_C is the blow up of \mathbb{P}^2 at 3 non collinear points and that these 3 exceptional divisors are the only curves of M_C with bidegree $(1, 0)$. Since M_C has only finitely many curves of bidegree $(1, 0)$, D is the same for all non-zero elements of \mathcal{V} and hence $\dim \mathcal{V} = 1$. \square

The following result completes the proof of Theorem 1.3.

Theorem 4.10. *There exist irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ containing exactly 4 twistor fibers and for any such S and $A \in \mathcal{T}(4)$ with $A \subset S$, there exists a curve L of bidegree $(1, 0)$ intersecting each connected component of A . Furthermore, $h^0(J_A(1, 2)) = 2$ and each $S \in |J_A(1, 2)|$ is singular along L .*

Proof. By Theorem 1.4 no irreducible surface of bidegree $(1, 2)$ contains at least 5 twistor fiber. The curve L exists by Lemma 4.8. Now we reverse the construction. We start with $A \in \mathcal{T}(4, L)^-$. Let $2L$ be the closed subscheme of the “double line”. To prove that every $S \in |J_A(1, 2)|$ is singular at every point of L , it suffices to prove that $h^0(J_A(1, 2)) = h^0(J_{A \cup 2L}(1, 2))$. Lemma 4.9 gives $h^0(J_A(1, 2)) = 2$. Hence, it is sufficient to prove that $h^0(J_{A \cup 2L}(1, 2)) > 1$. For

any connected component C of A , let M_C be the only surface of bidegree $(1, 1)$ containing $A \setminus C$ and let Y_C be the only surface of bidegree $(0, 1)$. Since $C \cap L \neq \emptyset$, $L \cap Y_C \neq \emptyset$. Since Y_C has bidegree $(0, 1)$ and L bidegree $(1, 0)$, we get $L \subset Y_C$. Thus $L \subseteq M_C \cap Y_C$ and hence $|\mathcal{J}_{A \cup 2L}(1, 2)|$ contains at least the 4 reducible elements of $|\mathcal{J}_A(1, 2)|$. Hence $h^0(\mathcal{J}_{A \cup 2L}(1, 2)) > 1$. \square

4.2. Non existence results for surfaces of bidegree $(1, 2)$ and $(1, 3)$. In this last section, we prove our last two main results, i.e. Theorems 1.4 and 1.5.

For any $A \in \mathcal{T}(n)^-$, $n \geq 4$, let us call L and $R := j(L)$ the curves of bidegree $(1, 0)$ e $(0, 1)$ respectively, intersecting all the connected components of A .

In view of our goal, we need to discuss the reducibility of some surfaces containing a certain number of twistor fibers. First of all, fix an integer $n \geq 2$, take $B \in \mathcal{T}(4)$ such that $h^0(\mathcal{J}_B(1, 1)) > 0$ and call M the unique (see e.g. Remark 4.2) surface of bidegree $(1, 1)$ containing B . Since every element of $\mathcal{C}(1)$ is contained in an element of $|\mathcal{O}_{\mathbb{F}}(0, 1)|$ for each $E \in \mathcal{T}(n - 1)$ there exists a reducible element $W \in |\mathcal{O}_{\mathbb{F}}(1, k)|$, union of M and $n - 1$ surfaces of bidegree $(0, 1)$ such that $B \cup E \subset W$. The following lemma is a kind of inverse of this remark. Moreover, it will also be a key tool in the last two proofs.

Lemma 4.11. *If $d \geq 2$ and $A \in \mathcal{T}(d + 3)^-$ are such that $h^0(\mathcal{J}_A(1, d)) > 0$, then every element of $|\mathcal{J}_A(1, d)|$ has an irreducible component M of bidegree $(1, 1)$ containing at least 4 connected components of A . In particular, for any $n \geq d + 3$, there is no irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, d)|$ containing $A \in \mathcal{T}(n)^-$.*

Proof. In order to prove the last statement, it is sufficient to do the case $n = d + 3$, and thus it is sufficient to prove the first assertion.

We use induction on $d \geq 2$. Let us first assume $d = 2$. Take $A \in \mathcal{T}(5)^-$ and let L and $j(L)$ be the curves of bidegree $(1, 0)$ and $(0, 1)$ intersecting all the connected components of A . Fix a connected component C of A and set $B := A \setminus C$. Since $C \cap L \neq \emptyset$, the curve $C \cup L$ is a connected and nodal curve of bidegree $(2, 1)$ with arithmetic genus 0. Hence $h^0(\mathcal{O}_{C \cup L}(0, 1)) = 2$. Thus there is $Y \in |\mathcal{J}_{C \cup L}(0, 1)|$ and such a Y is unique. Since any two smooth conics of Y meet, no component of B is contained in Y . Therefore, $B \cap Y$ is formed by 4 points of $L \setminus (L \cap C)$. Recall that $\mathcal{O}_Y(1, 2) \cong \mathcal{O}_{F_1}(h + 3f)$ and that $C \in |\mathcal{O}_{F_1}(h + f)|$ and thus $\mathcal{J}_{A \cap Y, Y} \cong \mathcal{J}_{B \cap L, Y}(2f)$. Since every element of $|f|$ contains a unique point of L we have that $h^0(D, \mathcal{J}_{A \cap Y, Y}(1, 2)) = 0$. The residual exact sequence of Y

$$0 \rightarrow \mathcal{J}_B(1, 1) \rightarrow \mathcal{J}_A(1, 2) \rightarrow \mathcal{J}_{A \cap Y, Y}(1, 2) \rightarrow 0,$$

gives an isomorphism $\varphi : H^0(\mathcal{J}_B(1, 1)) \rightarrow H^0(\mathcal{J}_A(1, 2))$. If $h^0(\mathcal{J}_B(1, 1)) = 0$, then $h^0(\mathcal{J}_A(1, 2)) = 0$. Now suppose $h^0(\mathcal{J}_B(1, 1)) \neq 0$. The isomorphism φ says that every $W \in |\mathcal{J}_A(1, 2)|$ has Y as an irreducible component, say $W = Y \cup W_1$ with $W_1 \in |\mathcal{J}_B(1, 1)|$, and hence we have the thesis.

Now assume $d \geq 3$ and use induction on d . By reasoning as in the base case, take $A \in \mathcal{T}(d + 3)^-$ and use the exact sequence

$$0 \rightarrow \mathcal{J}_B(1, d - 1) \rightarrow \mathcal{J}_A(1, d) \rightarrow \mathcal{J}_{A \cap Y, Y}(1, d) \rightarrow 0,$$

to prove that $h^0(Y, \mathcal{J}_{A \cap Y, Y}(1, d)) = 0$ and thus that there is an isomorphism $\varphi : H^0(\mathcal{J}_B(1, d - 1)) \rightarrow H^0(\mathcal{J}_A(1, d))$. Now, again, if $h^0(\mathcal{J}_B(1, d - 1)) = 0$, then $h^0(\mathcal{J}_A(1, d)) = 0$. So we assume $h^0(\mathcal{J}_B(1, d - 1)) \neq 0$. The isomorphism φ says that each $S \in |\mathcal{J}_A(1, d)|$ has Y as an irreducible component, i.e. $S = D \cup S_1$ with $S_1 \in |\mathcal{J}_B(1, d - 1)|$. The inductive assumption says that S_1 has an irreducible component M of bidegree $(1, 1)$ containing at least 4 components of B . \square

We now have all the ingredients to prove Theorems 1.4 and 1.5. First, we prove that no irreducible surface of bidegree $(1, 2)$ contains 5 twistor fibers.

Proof of Theorem 1.4. Suppose there exists an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, 2)|$ containing $A \in \mathcal{T}(5)$. Lemma 4.8 shows that for any union $A' \subset A$ of 4 components of A , there exists a union $A'' \subset A'$ of 3 connected components intersecting some L of bidegree $(1, 0)$. Let L be a curve of bidegree $(1, 0)$ intersecting the maximal number, z , of components of A . Obviously $z \geq 3$. By Lemma 4.11 to get a contradiction, it is sufficient to prove that $z \geq 5$.

Assume $z \in \{3, 4\}$. Take any order C_1, \dots, C_5 of the connected components of A and set $B_i := \pi_1(C_i)$, $1 \leq i \leq 5$. Each B_i is a line of \mathbb{P}^2 . Since any two conics contained in an element of $|\mathcal{O}_{\mathbb{F}}(1, 0)|$ meet, B_1, \dots, B_5 are 5 different lines of \mathbb{P}^2 . For any $i < j < h$ there exists a curve T of bidegree $(1, 0)$ intersecting C_i, C_j and C_h if and only if B_h contains the point $B_i \cap B_j$ and in this case $L = \pi_1^{-1}(C_i \cap C_j)$. Without loss of generality, we can assume that L meets C_1, \dots, C_z .

(a) Assume $z = 3$ and hence $B_1 \cap B_2 \in B_3$. Applying Lemma 4.8 to $C_1 \cup C_2 \cup C_4 \cup C_5$ we have one of the following mutually exclusive relations:

$$\begin{aligned} B_1 \cap B_2 \cap B_4 &\neq \emptyset, & B_1 \cap B_2 \cap B_5 &\neq \emptyset, \\ B_1 \cap B_4 \cap B_5 &\neq \emptyset, & B_2 \cap B_4 \cap B_5 &\neq \emptyset. \end{aligned}$$

Since $B_1 \cap B_2 \in B_3$ and $z = 3$, we can exclude the first two cases, i.e. we have

$$B_1 \cap B_2 \cap B_4 = B_1 \cap B_2 \cap B_5 = \emptyset.$$

Thus, either $B_1 \cap B_4 \cap B_5 \neq \emptyset$ or $B_2 \cap B_4 \cap B_5 \neq \emptyset$. Exchanging if necessary C_1 and C_2 we may assume $B_1 \cap B_4 \cap B_5 \neq \emptyset$, i.e. $B_4 \cap B_5 \in B_1$, and hence $B_2 \cap B_4 \cap B_5 = \emptyset$. Since $B_2 \cap B_4 \cap B_5 = \emptyset$, applying Lemma 4.8 to $C_2 \cup C_3 \cup C_4 \cup C_5$ we have one of the following mutually exclusive relations

$$\begin{aligned} B_2 \cap B_3 \cap B_4 &\neq \emptyset, & B_2 \cap B_3 \cap B_5 &\neq \emptyset, \\ B_3 \cap B_4 \cap B_5 &\neq \emptyset. \end{aligned}$$

Since $B_4 \cap B_5 \in B_1$ and $z = 3$, $B_3 \cap B_4 \cap B_5 = \emptyset$. Since $B_2 \cap B_3 \in B_1$ and $z = 3$, $B_2 \cap B_3 \cap B_5 = B_2 \cap B_3 \cap B_4 = \emptyset$, a contradiction.

(b) Assume $z = 4$. Since $C_1 \cap L \neq \emptyset$, the curve $C_1 \cup L$ is a connected and nodal curve of arithmetic genus 0 and bidegree $(2, 1)$. Thus $h^0(\mathcal{O}_{C_1 \cup L}(0, 1)) = 2$. Thus there is $Y \in |\mathcal{J}_{C_1 \cup L}(0, 1)|$. Since S is irreducible, Y is not an irreducible component of S . So the residual exact sequence of Y

$$0 \rightarrow \mathcal{J}_B(1, 1) \rightarrow \mathcal{J}_A(1, 2) \rightarrow \mathcal{J}_{A \cap Y, Y}(1, 2) \rightarrow 0,$$

gives $h^0(Y, \mathcal{J}_{A \cap Y, Y}(1, 2)) \neq 0$. Up to the isomorphism of Y and F_1 we have $L = h, C_1 \in |\mathcal{O}_{F_1}(h + f)|$ and $\mathcal{O}_Y(1, 2) \cong \mathcal{O}_{F_1}(h + 3f)$. Since $(A \setminus C_1) \cap C_1 = \emptyset$, $\mathcal{J}_{A \cap Y, Y}(1, 3) \cong \mathcal{J}_{(A \setminus C_1) \cap D, D}(2f)$. Since L meets C_1, \dots, C_4 , $(A \setminus C_1) \cap D$ contains a set $F \subset L$ such that $\#F = 3$. Since every element of $|f|$ contains a unique point of L , $h^0(Y, \mathcal{J}_{A \setminus C \cap Y, Y}(2f)) = 0$, a contradiction. \square

We now conclude with the proof of Theorem 1.5, which concerns surfaces of bidegree $(1, 3)$.

Proof of Theorem 1.5: Assume the existence of $A \in \mathcal{T}(6)$ and of an irreducible $S \in |\mathcal{O}_{\mathbb{F}}(1, 3)|$ containing A . By Lemma 4.11 to get a contradiction it is sufficient to prove the existence of a curve L of bidegree $(1, 0)$ such that all the components of A intersect L . By Lemma 4.7 for any union $A' \subset A$ of 5 components of A there is a union $A'' \subset A'$ of 3 connected components intersecting some L of bidegree $(1, 0)$. Let L be a curve of bidegree $(1, 0)$ intersecting the maximal number, z , of components of A . We have that $z \geq 3$. Therefore, by Lemma 4.11 it is sufficient to prove that $z \geq 6$. Assume then that $z \leq 5$. We will now exclude all the cases $z = 3, 4, 5$.

For any connected component C of A , Lemma 4.7 tells us that there exists a curve L of bidegree $(1, 0)$ which intersects at least 3 connected components of $A \setminus C$. In particular there is an irreducible $M \in |\mathcal{O}_{\mathbb{F}}(1, 1)|$ containing at least 3 components of A (see Remark 4.2). Note that $j(M) = M$. We can take M with the additional condition that it contains the maximal number e of components of A . Let E be the union of the components of A contained in M . So $3 \leq e \leq z \leq 5$.

Since each twistor fiber is j -invariant, $j(L)$ hits every connected component of E . Bézout gives $L \cup j(L) \subset M$ and $L \subset S$. If $e \geq 4$, Bézout gives $j(L) \subset S$. However, the one-dimensional cycle $M \cap S$ has bidegree $(7, 5)$ and thus $e \leq 4$. Set $\Sigma := S \cap M$ (as a scheme-theoretic intersection). Since the one-dimensional scheme Σ is the complete intersection of \mathbb{F} with 2 very ample divisors, $h^0(\mathcal{O}_{\Sigma}) = 1$. Set $F := A \setminus E$.

(a) Assume $e = 4$. So $E \cup L \cup j(L) \subset \Sigma$. Since $E \cup L \cup j(L)$ has bidegree $(5, 5)$ and $h^0(\mathcal{O}_{\Sigma}) = 1$, Σ is the union of $E \cup L \cup j(L)$ and a multiple structure on L . Note that $\Sigma \in |\mathcal{O}_{\mathbb{F}}(1, 3)|$ and that Σ contains $E \cup j(L)$ with multiplicity 1 and L with multiplicity 3 (as divisors of the smooth surface M). Since Σ has multidegree $(7, 5)$, $\Sigma = 3L \cup j(L) \cup E$. Note that Σ contains the degree 4 zero-dimensional scheme $F \cap M$. Since $F \cap E = \emptyset$, $F \cap (j(L) \cup L) \neq \emptyset$. Thus at least one irreducible component, T , of F meets $L \cup j(L)$. Since $j(T) = T$, $T \cap L \neq \emptyset$. Thus $z = 5$. Let C be a component of E . Since $C \cap L \neq \emptyset$, $C \cup L$ is a connected and nodal curve of bidegree $(2, 1)$ with arithmetic genus 0. Thus $h^0(\mathcal{O}_{C \cup L}(0, 1)) = 2$. So there is $Y \in |\mathcal{J}_{C \cup L}(0, 1)| \neq \emptyset$. Since S is irreducible, Y is not an irreducible component of S . Thus the residual exact sequence of Y gives $h^0(Y, \mathcal{J}_{A \cap Y, Y}(1, 3)) \neq 0$. Up to the isomorphism of Y and F_1 we have $L = h, C \in |\mathcal{O}_{F_1}(h + f)|$ and $\mathcal{O}_Y(1, 3) \cong \mathcal{O}_{F_1}(h + 4f)$. Since $(A \setminus C) \cap C = \emptyset$, $\mathcal{J}_{A \cap Y, Y}(1, 3) \cong \mathcal{J}_{(A \setminus C) \cap Y, Y}(3f)$. Since $z = 5$, $(A \setminus C) \cap Y$ contains a set $H \subset L$

such that $\#H = 4$. Since every element of $|f|$ contains a unique point of L , $h^0(Y, \mathcal{J}_{A \setminus C} \cap Y, Y(3f)) = 0$, a contradiction.

(b) Assume $e = 3$. Fix a connected component C of E and set $B := A \setminus C$. Set $\{Y\} := |\mathcal{J}_C(0, 1)|$. As in step (a), we have that $L \subset Y$. The following exact sequence

$$0 \rightarrow \mathcal{J}_B(1, 2) \rightarrow \mathcal{J}_A(1, 3) \rightarrow \mathcal{J}_{C \cup (B \cap Y), Y}(1, 3) \rightarrow 0 \quad (23)$$

is the residual exact sequence of Y . Since Y is not an irreducible component of S , we have $h^0(Y, \mathcal{J}_{C \cup (B \cap Y), Y}(1, 3)) > 0$. As in step (a), we have $\mathcal{J}_{C \cup (B \cap Y), Y}(1, 3) \cong \mathcal{J}_{B \cap Y, F_1}(3f)$. We now have two possibilities: $h^0(\mathcal{J}_B(1, 2)) = 0$ or $h^0(\mathcal{J}_B(1, 2)) > 0$.

(b1) Let us assume for the moment that $h^0(\mathcal{J}_B(1, 2)) = 0$. Then we have $h^0(Y, \mathcal{J}_{C \cup (B \cap Y), Y}(1, 3)) \geq 2$ and $h^1(Y, \mathcal{J}_{C \cup (B \cap Y), Y}) \geq 3$. Up to the identification of Y and F_1 we have $\mathcal{J}_{C, Y}(1, 3) \cong \mathcal{O}_{F_1}(3f)$. So the 5 points $B \cap Y$ give at most one condition to the linear system $|\mathcal{O}_{F_1}(3f)|$. Thus there is $J \in |\mathcal{O}_{F_1}(f)|$ such that $B \cap Y \subset J$. Note that J is a curve of bidegree $(1, 0)$. The maximality of the integer e gives a contradiction.

(b2) Assume that $h^0(\mathcal{J}_B(1, 2)) > 0$. By Theorem 1.4 every surface containing B is reducible, say $M_1 \cup Y$ with M_1 irreducible of bidegree $(1, 1)$ containing at least 4 components of B . Thus $e \geq 4$, a contradiction. \square

References

- [1] ALTAVILLA, AMEDEO; BALLICO, EDOARDO. Twistor lines on algebraic surfaces. *Ann. Global Anal. Geom.* **55** (2019), no. 3, 555–573. [MR3936233](#), [Zbl 07050118](#), doi: [10.1007/s10455-018-9640-2](#). [1117](#)
- [2] ALTAVILLA, AMEDEO; SARFATTI, GIULIA. Slice-polynomial functions and twistor geometry of ruled surfaces in $\mathbb{C}\mathbb{P}^3$. *Math. Z.* **291** (2019), no. 3-4, 1059–1092. [MR3936099](#), [Zbl 1415.53035](#), doi: [10.1007/s00209-018-2225-8](#). [1117](#)
- [3] ALTAVILLA, AMEDEO; BALLICO, EDOARDO; BRAMBILLA, MARIA C. Surfaces in the flag threefold containing smooth conics and twistor fibers. *Mediterr. J. Math.* **19** (2022). [MR4508025](#), [Zbl 1510.32043](#), doi: [10.1007/s00009-022-02202-3](#). [1118](#), [1121](#), [1123](#), [1124](#), [1125](#), [1126](#), [1128](#), [1139](#), [1141](#)
- [4] ALTAVILLA, AMEDEO; BALLICO, EDOARDO; BRAMBILLA, MARIA C.; SALAMON, SIMON. Twistor geometry of the Flag manifold. *Math. Z.* **303** (2023), no. 24. [MR4522734](#), [Zbl 1507.14069](#), doi: [10.1007/s00209-022-03161-x](#). [1118](#), [1119](#), [1121](#), [1122](#), [1123](#), [1124](#), [1129](#), [1130](#)
- [5] ARMSTRONG, JOHN; POVERO, MASSIMILIANO; SALAMON, SIMON. Twistor lines on cubic surfaces. *Rend. Semin. Mat. Univ. Politec. Torino* **71** (2013), no. 3-4, 317–338. [MR3506390](#), [Zbl 1332.53063](#). [1117](#), [1118](#)
- [6] ATIYAH, MICHAEL F.; HITCHIN, NIGEL J.; SINGER, ISADORE M. Self-duality in four-dimensional Riemannian geometry. *Proc. Roy. Soc. London Ser. A* **362** (1978), no. 1711, 425–461. [MR0506229](#), [Zbl 0389.53011](#). [1117](#), [1119](#)
- [7] CHIRKA, EVGENII M. Orthogonal complex structures in \mathbb{R}^4 . (Russian) *Uspekhi Mat. Nauk* **73** (2018), no. 1, 99–172. Translation in *Russian Math. Surveys* **73** (2018), no. 1, 91–159. [MR3749619](#), [Zbl 1400.32013](#), doi: [10.4213/rm9788](#). [1117](#)
- [8] FUJIKI, AKIRA; PONTECORVO, MASSIMILIANO. Twistors and bi-Hermitian surfaces of non-Kähler type. *SIGMA Symmetry Integrability Geom. Methods Appl.* **10** (2014), Paper 042, 13 pp. [MR3210593](#), [Zbl 1288.53038](#), doi: [10.3842/SIGMA.2014.042](#). [1117](#)

- [9] GENTILI, GRAZIANO; SALAMON, SIMON; STOPPATO, CATERINA. Twistor transforms of quaternionic functions and orthogonal complex structures. *J. Eur. Math. Soc. (JEMS)* **16** (2014), no.11, 2323?2353. [MR3283400](#), [Zbl 1310.53045](#), doi: [10.4171/JEMS/488](#). [1117](#)
- [10] HARTSHORNE, ROBIN Algebraic Geometry. *Springer-Verlag, Berlin–Heidelberg–New York*, 1977. xvi+496 pp. [MR0463157](#), [Zbl 0367.14001](#).
- [11] HITCHIN NIGEL. Kählerian twistor spaces. *Proc. London Math. Soc. (3)* **43** (1981), no.1, 133–150. [MR0623721](#), [Zbl 0474.14024](#), doi: [10.1112/plms/s3-43.1.133](#). [1117](#), [1118](#), [1119](#)
- [12] PONTECORVO, MASSIMILIANO. Complex structures on Riemannian four-manifolds. *Math. Ann.* **309** (1997), no.1, 159–177. [MR1467652](#), [Zbl 0893.53026](#), doi: [10.1007/s002080050108](#). [1117](#)
- [13] SALAMON, SIMON; VIACLOVSKY, JEFF. Orthogonal complex structures on domains in \mathbb{R}^4 . *Math. Ann.* **343** (2009), no.4, 853–899. [MR2471604](#), [Zbl 1167.32017](#), doi: [10.1007/s00208-008-0293-5](#). [1117](#), [1118](#)

(Amedeo Altavilla) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI BARI ‘ALDO MORO’, VIA EDOARDO ORABONA, 4, 70125, BARI, ITALIA
amedeo.altavilla@uniba.it

(Edoardo Ballico) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TRENTO, VIA SOMMARIVE 14, 38123, POVO, TRENTO, ITALIA
edoardo.ballico@unitn.it

(Maria Chiara Brambilla) UNIVERSITÀ POLITECNICA DELLE MARCHE, VIA BRECCE BIANCHE, I-60131 ANCONA, ITALIA
m.c.brambilla@univpm.it

This paper is available via <http://nyjm.albany.edu/j/2023/29-45.html>.