

An arithmetic Kontsevich–Zorich monodromy of a symmetric origami in genus 4

Xun Gong and Anthony Sanchez

ABSTRACT. We demonstrate the existence of a certain genus four origami whose Kontsevich–Zorich monodromy is arithmetic in the sense of Sarnak. The surface is interesting because its Veech group is as large as possible and given by $SL(2, \mathbb{Z})$. When compared to other surfaces with Veech group $SL(2, \mathbb{Z})$ such as the Eierlegendre Wollmichsau and the Ornithorynque, an arithmetic Kontsevich–Zorich monodromy is surprising and indicates that there is little relationship between the Veech group and monodromy group of origamis. Additionally, we record the index and congruence level in the ambient symplectic group which gives data on what can appear in genus 4.

CONTENTS

1. Introduction	1413
2. The origami \mathcal{O}	1416
3. Homology of \mathcal{O}	1417
4. Kontsevich–Zorich monodromy group of \mathcal{O}	1419
5. Zariski denseness, arithmeticity, index, and congruence level	1420
References	1423

1. Introduction

The Kontsevich–Zorich monodromy group encodes the homological data of translation surfaces along $SL(2, \mathbb{R})$ -orbits. Due to the intersection form on the underlying surface, the Kontsevich–Zorich monodromy group lies as a subgroup of a symplectic group and creates a natural bridge between the geometric objects of translation surfaces and the Kontsevich–Zorich monodromy group and its algebraic nature. One such question that has received much investigation is whether the Kontsevich–Zorich monodromy group is arithmetic or is thin in the sense of Sarnak [15]. An origami has *arithmetic Kontsevich–Zorich monodromy* if the Kontsevich–Zorich monodromy group has finite index in the

Received August 4, 2023.

2020 *Mathematics Subject Classification*. Primary 37D40; Secondary 32G15.

Key words and phrases. Translation surfaces, Kontsevich–Zorich monodromy group, origami, arithmetic monodromy group.




FIGURE 1. The origami \mathcal{O} with identification labeled by a permutation.

integer matrices of its Zariski closure and *thin* if the Kontsevich–Zorich monodromy group has infinite index in the integer matrices of its Zariski closure.

In genus 2, the arithmeticity of all origami was established by Möller, see also Appendix B of [1] for a proof. In genus 3, an origami with arithmetic monodromy in the minimal stratum was established in Hubert–Matheus [9]. Shortly thereafter, an infinite family of origami in genera 3, 4, 5, and 6 with arithmetic monodromy were exhibited in [1, 10].

Assuming arithmeticity has been established, finer questions regarding the algebraic nature of the monodromy group, such as the index and what type of group is obtained, have been considered. Experimental data [1] shows that for genus 2 origamis with two conical singularities and less than 22 squares, the index is 1, 3, 4, 6, 12, or 24. For genus 2 origami with a single conical singularity, Kattler [11] has shown part of a conjecture that the index is always 1 or 3. Concerning what type of groups are obtained, in the context of symplectic groups, there is a positive solution to the congruence subgroup property. That is, all finite index subgroups of symplectic groups, such as the Kontsevich–Zorich monodromy groups in the works mentioned above, are a congruence group of some level l . In [1], they show a specific origami in genus 3 is conjugate, up to finite index, to a congruence subgroup of level 16 and of index 46080 in $\mathrm{Sp}(4, \mathbb{Z})$.

Our main result is a new addition to the list of origami with arithmetic monodromy. Later in the introduction, we will see that this surface being arithmetic is surprising and relate this to the monodromy of other surfaces. Additionally, we record the first index and congruence level of a surface in genus 4 which yields experimental data for what can occur in the context of genus 4 origami. See Figure 1 for an image of the surface.

Theorem 1.1. *The non-tautological part of the Kontsevich–Zorich monodromy group Γ associated to a certain origami \mathcal{O} of genus 4 is arithmetic. More precisely,*

up to a finite index subgroup that conjugates Γ to a subgroup of the standard symplectic group $\mathrm{Sp}(6, \mathbb{Z})$, the Kontsevich–Zorich monodromy group is a congruence subgroup of level 64 and of index $4156153952993280 = 2^{42} \cdot 3^3 \cdot 5 \cdot 7$.

As we will see in Section 3, the intersection matrix of the homology we use does not have determinant one, so we must pass to a finite index subgroup of Γ in order to be a subgroup of the standard symplectic group. The index and congruence level are computed through GAP packages developed by Detinko–Flannery–Hulpke [3] and Kattler–Weitze–Schmithüsen [12]. Theorem 1.1 results in the first data point of the possible indices and congruence levels of origami in genus 4.

Relation to other results. While an infinite family of arithmetic monodromies in genus 4 was established by Kany–Matheus [10], we believe the results of this article are worthwhile because of the unique properties of the surface. What drew our attention to this origami was that it is “exceptionally symmetric” in that its Veech group is as large as possible and given by $\mathrm{SL}(2, \mathbb{Z})$. Additionally, the homological dimension is always 2. We now give some context as to why these points are interesting.

It was shown experimentally in Shrestha–Wang [16], that there are only 4 origamis of less than 17 squares whose Veech group is $\mathrm{SL}(2, \mathbb{Z})$: the torus, the *Eierlegendre Wollmichsau*, the *Ornithorynque*, and the surface considered in this paper. We refer to [16] for the description of the Eierlegendre Wollmichsau and Ornithorynque. These two surfaces have been the focus of much study [14, 8, 5] because of their peculiar properties e.g. their Veech group is $\mathrm{SL}(2, \mathbb{Z})$ and they have totally degenerate Lyapunov spectrum. What’s most relevant for the discussion here is that Matheus–Yoccoz [14] showed the Kontsevich–Zorich monodromy groups of these two surfaces are finite groups. That the surface we consider in this article is arithmetic and thus “large” is quite interesting given that the monodromies of the Eierlegendre Wollmichsau and Ornithorynque are at the other end of the spectrum. This indicates that there is little relationship between the symmetries of the surface (the Veech group) and the Kontsevich–Zorich monodromy group of the surface.

Another interesting aspect of our surface is that the homological dimension is always two. Surfaces with directions of homological dimension one have that their action by the affine diffeomorphisms on the non-tautological part of homology is never faithful since Dehn multitwists in those directions produce elements in the kernel of the action and faithfulness (along with Zariski density) is enough for thinness (see Section 4.3 of [9]). Thus, our surface was a candidate to produce an origami with thin monodromy.

Organization. As in [9], we assume some familiarity with the basic features of origamis including their representation theory, homology, and Veech and affine groups of origamis. We recommend the survey of Matheus [13]. The rest of the note is dedicated to proving Theorem 1.1. The organization of the paper is as follows. In Section 2, we describe some properties of the origami we are studying. In Section 3, we find a basis for the homology of our origami. In

Section 4, we compute generators for the monodromy group. Lastly, in Section 5, we utilize an arithmeticity criterion of [17] that was explicated in [1] in the context of monodromies of origamis to prove Theorem 1.1.

1.1. Acknowledgements. The authors would like to thank Carlos Matheus Silva Santos and Pascal Kattler for a careful reading of the manuscript. In particular, we thank Matheus for comments on the Lyapunov spectrum and the reference [7] and Pascal Kattler for pointing out a mistake in a previous computation of $\alpha(S)$. Lastly, we are immensely grateful to Pascal Kattler and Gabriela Weitze-Schmithüsen for running the generators of Theorem 4.1 through their software [12]. A.S. was supported by the National Science Foundation Postdoctoral Fellowship under grant number DMS-2103136.

The authors are grateful to the anonymous referee for a careful reading of the manuscript and many useful suggestions.

2. The origami \mathcal{O}

Let \mathcal{O} denote the origami (see Figure 1 for an image of the surface) defined by the permutations

$$h = (1, 2, 3, 4, 5, 6)(12, 11, 10, 9, 8, 7)(13, 14)(15, 16)$$

and

$$v = (12, 2, 16, 14, 10, 6)(11, 5, 15, 13, 7, 1)(3, 9)(4, 8).$$

The commutator

$$[h, v] := v h v^{-1} h^{-1} = (1, 3, 5)(2)(4)(6)(7, 15, 9)(8)(10, 16, 12)(11)(13)(14)$$

has 3 non-trivial cycles of length 3 and so $\mathcal{O} \in \mathcal{H}(2, 2, 2)$ and has genus 4.

As mentioned in the introduction, one aspect that makes this surface so special is its Veech group. The Veech group is the stabilizer of \mathcal{O} with respect to the $\mathrm{SL}(2, \mathbb{R})$ action on the space of translation surfaces. let $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $S = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ and recall that $\mathrm{SL}(2, \mathbb{Z})$ is generated by these two matrices.

Proposition 2.1. *The Veech group of \mathcal{O} is $\mathrm{SL}(2, \mathbb{Z})$.*

Proof. At the level of permutations, the generators of $\mathrm{SL}(2, \mathbb{Z})$ act by the rules $T(h, v) = (h, v h^{-1})$ and $S(h, v) = (h v^{-1}, v)$. Since we care about the surfaces themselves rather than the labeling of the squares, the pair (h, v) is defined up to simultaneous conjugations i.e. (h, v) and $(\psi h \psi^{-1}, \psi v \psi^{-1})$ are regarded as the same origami. With this in mind, we let

$$\psi = (1, 3, 5)(2, 4, 6)(7, 10)(8, 11)(9, 12)(13)(14)(15, 16)$$

and

$$\phi = (1, 15)(2, 14, 6)(3, 9)(4)(5, 7)(8)(10, 12, 16)(11, 13).$$

A computation shows that $\psi \circ T(h, v) \circ \psi^{-1} = (h, v)$ and $\phi \circ S(h, v) \circ \phi^{-1} = (h, v)$. \square




FIGURE 2. Basis for the homology of \mathcal{O}

Remark 2.2. We remark that the work of Herrlich [7] yields that any origami can be covered by one whose Veech group is $SL(2, \mathbb{Z})$. However, explicit computations outside of the present article seem difficult because the genus of the cover can grow quickly. For example, for the L-shaped origami of genus 2 and with 3 squares, the corresponding cover is of genus 37 and has 108 squares.

In the introduction, we also claimed that the homological dimension is always two. That is, in any rational direction, the surface \mathcal{O} decomposes into cylinders whose waist curves span a two-dimensional subspace of the homology. Such surfaces have the potential for a thin monodromy group.

Proposition 2.3. *The homological dimension of \mathcal{O} in any rational direction is 2.*

Proof. The surface \mathcal{O} has two horizontal cylinders of length six and two horizontal cylinders of length two. Since the waist curves on the horizontal cylinders of length six disconnect \mathcal{O} , then they are homotopic. Similarly, the waist curves on the horizontal cylinders of length two are homotopic. By noting that the long and short waist curves are not homotopic, then we deduce that the homological dimension in the horizontal direction is two. Lastly, since the Veech group of \mathcal{O} is $SL(2, \mathbb{Z})$ and any other cylinder direction can be moved to the horizontal, then we conclude the homological direction in any rational direction is two. \square

Additionally, the homological dimension of \mathcal{O} has consequences on the Lyapunov spectrum of the Kontsevich-Zorich cocycle over $SL(2, \mathbb{R}) \cdot \mathcal{O}$. By Forni’s theorem [4], the second Lyapunov exponent is positive.

3. Homology of \mathcal{O}

In this section, we give a basis for the homology of \mathcal{O} and define certain subspaces needed for later sections. We will work with the following basis of the absolute homology group $H_1(\mathcal{O}, \mathbb{R})$ pictured in Figure 2 that we denote by $\{\gamma_1, \dots, \gamma_8\}$. The colors, orientations, and slopes are indicated in Table 1.

curve	color	slope	holonomy
γ_1	yellow	0	$(2, 0)^T$
γ_2	black	1	$(2, 2)^T$
γ_3	orange	0	$(6, 0)^T$
γ_4	red	1/3	$(18, 6)^T$
γ_5	grey	∞	$(0, 2)^T$
γ_6	light blue	-1	$(6, -6)^T$
γ_7	blue	∞	$(0, 6)^T$
γ_8	green	3	$(2, 6)^T$

TABLE 1. Some properties of a basis of the absolute homology of \mathcal{O} .

The corresponding intersection matrix of our basis of homology is:

$$\Omega = \begin{matrix} & \begin{matrix} \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 & \gamma_7 & \gamma_8 \end{matrix} \\ \begin{matrix} \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \gamma_4 \\ \gamma_5 \\ \gamma_6 \\ \gamma_7 \\ \gamma_8 \end{matrix} & \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & -1 & 1 & 1 \\ 0 & 0 & -1 & -1 & 0 & -1 & 1 & 1 \\ 0 & 1 & 0 & 2 & 1 & -2 & 2 & 2 \\ -1 & 1 & -2 & 0 & 2 & -9 & 7 & 5 \\ 0 & 0 & -1 & -2 & 0 & -1 & 0 & 0 \\ 1 & 1 & 2 & 9 & 1 & 0 & 2 & 3 \\ -1 & -1 & -2 & -7 & 0 & -2 & 0 & -1 \\ -1 & -1 & -2 & -5 & 0 & -3 & 1 & 0 \end{pmatrix} \end{matrix}.$$

Each entry of the matrix, $\Omega_{i,j}$, represents the intersection form $\langle \gamma_i, \gamma_j \rangle$, $i, j \in \{1, 2, \dots, 8\}$. The determinant of Ω is 16.

There is always the decomposition of the absolute homology

$$H_1(\mathcal{O}) = H_1^{\text{st}}(\mathcal{O}) \oplus H_1^{(0)}(\mathcal{O}),$$

where $H_1^{\text{st}}(\mathcal{O})$ the *tautological plane* spanned by the real and imaginary parts of the implicit Abelian differential and $H_1^{(0)}(\mathcal{O})$ is the 6-dimensional orthogonal complement (with respect to the intersection form) given by the *zero-holonomy subspace*. The name “zero-holonomy” comes from the fact that the horizontal and vertical displacement (with respect to the singular flat metric) of the curves are zero. Next section we justify the terminology of “tautological plane”. In the setting of our specific surface, it can be checked that with respect to the basis $\{\gamma_1, \dots, \gamma_8\}$ we have

$$H_1^{\text{st}}(\mathcal{O}) = \{2(\gamma_1 + \gamma_3), 2(\gamma_5 + \gamma_7)\} \text{ and } H_1^{(0)}(\mathcal{O}) = \{\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4, \epsilon_5, \epsilon_6\}$$

where

$$\begin{aligned} \epsilon_1 &= -3\gamma_1 + \gamma_3, & \epsilon_2 &= -6\gamma_1 - 3\gamma_2 + \gamma_4, \\ \epsilon_3 &= \gamma_1 - \gamma_2 + \gamma_5, & \epsilon_4 &= -6\gamma_1 + 3\gamma_2 + \gamma_6, \\ \epsilon_5 &= 3\gamma_1 - 3\gamma_2 + \gamma_7, & \epsilon_6 &= 2\gamma_1 - 3\gamma_2 + \gamma_8. \end{aligned}$$

4. Kontsevich–Zorich monodromy group of \mathcal{O}

In this section, we compute the Kontsevich–Zorich monodromy group of \mathcal{O} . We define these terms and encourage the reader to see Matheus [13] for more details. Let $\tilde{\alpha} : \text{Aff}(\mathcal{O}) \rightarrow \text{Sp}(H_1(\mathcal{O}, \mathbb{R}))$ denote the representation arising from the action of the affine diffeomorphisms $\text{Aff}(\mathcal{O})$ on the absolute homology group $H_1(\mathcal{O}, \mathbb{R})$. By noting that the automorphism group of \mathcal{O} is trivial, we can (and do) identify $\text{Aff}(\mathcal{O})$ with the Veech group of \mathcal{O} .

The representation $\tilde{\alpha}$ arising from the homological action of $\text{Aff}(\mathcal{O})$ respects this decomposition. In fact, the name “tautological” comes from the fact that $\tilde{\alpha}(g)|_{H_1^g} = g$. Since the action of $\text{Aff}(\mathcal{O})$ is well understood on the tautological plane, we restrict our analysis of the action of $\text{Aff}(\mathcal{O})$ to the zero-holonomy subspace. Let $\alpha : \text{Aff}(\mathcal{O}) \rightarrow \text{Sp}(H_1^{(0)}(\mathcal{O}))$ denote this restriction. The *Kontsevich–Zorich monodromy group* of \mathcal{O} is defined to be $\alpha(\text{Aff}(\mathcal{O}))$.

Theorem 4.1. *The Kontsevich-Zorich monodromy group of \mathcal{O} is generated by the following two matrices:*

$$\alpha(T) = \begin{pmatrix} 1 & 11 & 2 & -6 & 7 & 5 \\ 0 & -4 & -1 & 3 & -3 & -2 \\ 0 & -2 & -1 & 3 & -2 & -3 \\ 0 & -3 & -1 & 3 & -3 & -3 \\ 0 & 2 & 0 & -1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix} \text{ and } \alpha(S) = \begin{pmatrix} 1 & 1 & -1 & 4 & -3 & -3 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 1 & 6 & 2 & -3 & 3 & 3 \\ 0 & 3 & 1 & -3 & 3 & 3 \\ 1 & 0 & -1 & 3 & -2 & -1 \\ 0 & 4 & 1 & -3 & 3 & 2 \end{pmatrix}.$$

Proof. Since $\text{Aff}(\mathcal{O}) = \text{SL}(2, \mathbb{Z}) = \langle T, S \rangle$, then it suffices to compute the image $\alpha(T)$ and $\alpha(S)$. To this end, we utilize the intersection form on the surface \mathcal{O} . See for example, the proof on page 12 of [6]. We first compute the homological action of $\text{SL}(2, \mathbb{Z})$ given by the representation $\tilde{\alpha}$ and then restrict it to $\alpha = \tilde{\alpha}|_{H_1^{(0)}}$.

We only compute $\tilde{\alpha}(T)$ and a similar argument works for $\tilde{\alpha}(S)$. Given a basis curve of the absolute homology of \mathcal{O} , we consider the intersection pattern of this curve after the application of $\tilde{\alpha}(T)$ against the basis $\{\gamma_1, \dots, \gamma_8\}$ of $H_1(\mathcal{O})$. By expressing $\tilde{\alpha}(T)$ as a linear combination of the basis and utilizing the intersection form, we create a linear system of equations that we can solve.

For example, we compute $\tilde{\alpha}(T)\gamma_2$. The image is shown in Figure 3. The intersection pattern of $\langle \tilde{\alpha}(T)\gamma_2, \gamma_i \rangle$ with the basis can be recorded with the vector $(0, -1, -1, 1, -2, 1, 0)$. We note that this requires using the relabeling map ψ from Section 2. Now writing $\tilde{\alpha}(T)\gamma_2 = \sum_{i=1}^8 a_i\gamma_i$ where $a_i \in \mathbb{R}$ and using the




FIGURE 3. γ_2 (left) and $\tilde{\alpha}(T)\gamma_2$ (right)

intersection matrix Ω we get

$$\langle \tilde{\alpha}(T)\gamma_2, \gamma_j \rangle = \left\langle \sum_{i=1}^8 a_i \gamma_i, \gamma_j \right\rangle = \sum_{i=1}^8 a_i \langle \gamma_i, \gamma_j \rangle = \sum_{i=1}^8 a_i \Omega_{i,j}.$$

Setting this equal to the vector $(0, -1, -1, 1, -2, 1, 0)$ and solving the linear system yields $\tilde{\alpha}(T)\gamma_2 = -4\gamma_1 - 2\gamma_3 + \gamma_4 + \gamma_5 + \gamma_6$. Doing this for all the basis curves and doing a similar computation for $\tilde{\alpha}(S)$ shows that the homological is given by:

$$\begin{aligned} \tilde{\alpha}(T)\gamma_1 &= \gamma_1, & \tilde{\alpha}(S)\gamma_1 &= \gamma_2 \\ \tilde{\alpha}(T)\gamma_2 &= -4\gamma_1 - 2\gamma_3 + \gamma_4 + \gamma_5 + \gamma_6, & \alpha(S)\gamma_2 &= 2\gamma_1 + \gamma_3 - \gamma_5 - \gamma_6 + \gamma_7 - \gamma_8 \\ \tilde{\alpha}(T)\gamma_3 &= \gamma_3, & \tilde{\alpha}(S)\gamma_3 &= \gamma_1 - \gamma_2 + \gamma_3 + \gamma_5 + \gamma_7 \\ \tilde{\alpha}(T)\gamma_4 &= 7\gamma_1 - \gamma_2 + 5\gamma_3 - \gamma_4 + \gamma_5 + 2\gamma_7, & \tilde{\alpha}(S)\gamma_4 &= 5\gamma_1 + 4\gamma_3 - \gamma_4 + 3\gamma_5 + 3\gamma_7 + \gamma_8, \\ \tilde{\alpha}(T)\gamma_5 &= \gamma_2, & \tilde{\alpha}(S)\gamma_5 &= \gamma_5 \\ \tilde{\alpha}(T)\gamma_6 &= -\gamma_7, & \tilde{\alpha}(S)\gamma_6 &= \gamma_3 \\ \tilde{\alpha}(T)\gamma_7 &= \gamma_1 - \gamma_2 + \gamma_3 + \gamma_5 + \gamma_7, & \tilde{\alpha}(S)\gamma_7 &= \gamma_7 \\ \tilde{\alpha}(T)\gamma_8 &= -\gamma_1 - \gamma_3 + \gamma_4 + \gamma_7 - \gamma_8, & \tilde{\alpha}(S)\gamma_8 &= \gamma_1 + \gamma_2 + 2\gamma_7 - \gamma_8. \end{aligned}$$

Finally, using that the action on the zero-holonomy subspace is just the restriction of the homological action, $\alpha = \tilde{\alpha}|_{H_1^{(0)}}$, we obtain the matrices in the statement of the theorem with respect to the basis $\{\epsilon_1, \dots, \epsilon_6\}$. \square

5. Zariski denseness, arithmeticity, index, and congruence level

In this section, we verify the Zariski density and arithmeticity of the monodromy group $\Gamma = \alpha(\text{Aff}(\mathcal{O})) = \langle \alpha(T), \alpha(S) \rangle$ as well as record the index and congruence level of the conjugate of a finite index subgroup of Γ . The Zariski density will be verified using a criterion of Kany-Matheus [10] (bottom of page 2). The arithmeticity will be done by using a criterion of Singh–Venkataramana [17] and explicated for origamis in [1] page 10. Their metacode uses 2-cylinder decompositions which do not exist for our surface, so we apply a slight modification, see Remark 5.2. The index and congruence level is computed using the computer software of Detinko–Flannery–Hulpke [3] and Kattler–Weitze-Schmithüsen [12].

Theorem 5.1. *The monodromy group Γ is Zariski dense in $\mathrm{Sp}(6, \mathbb{R})$.*

Proof. According to Kany-Matheus [10], it suffices to find two elements A and B in $\mathrm{Sp}(H_1^{(0)}(\mathcal{O}))$ such that:

- (1) The matrix A is a *Galois-pinching*. That is, a symplectic matrix whose characteristic polynomial is irreducible over \mathbb{Z} , splits over \mathbb{R} , and possesses the largest possible (hyperoctahedral) Galois group among reciprocal polynomials of degree 6;
- (2) A non-trivial unipotent element B such that $(B - \mathrm{Id})(H_1^{(0)}(\mathcal{O}))$ is not a Lagrangian subspace of $H_1^{(0)}(\mathcal{O})$.

Let $A = \alpha(S)\alpha(T)\alpha(S)\alpha(T)^{20}$ and $B = \alpha(T)^6$. The characteristic polynomial of A is

$$f_A(x) = 1 - 3x - 91x^2 - 262x^3 - 91x^4 - 3x^5 + x^6.$$

Note that this polynomial is reciprocal: $x^6 f_A(x^{-1}) = f_A(x)$. Additionally, it splits over \mathbb{R} and is irreducible over \mathbb{Z} . Moreover, the permutation group acts on a set of cardinality 6, the order of the Galois group is precisely $2^4 \cdot 3 = 48$ which matches with the order of the hyperoctahedral group. That gives us condition (1).

Now we turn our attention to $B = \alpha(T)^6$. The subspace $(B - \mathrm{Id})(H_1^{(0)}(\mathcal{O}))$ is 1 dimensional. Yet, if $(B - \mathrm{Id})(H_1^{(0)}(\mathcal{O}))$ is a Lagrangian subspace, then

$$\dim((B - \mathrm{Id})(H_1^{(0)}(\mathcal{O}))) = \frac{\dim(H_1^{(0)}(\mathcal{O}))}{2} = \frac{6}{2} = 3 \neq 1.$$

Therefore, we have satisfied condition (2).

Together with the criterion, the monodromy Γ is Zariski dense in $\mathrm{Sp}(6, \mathbb{R})$. □

The Zariski-density provided by the above theorem, allows us to say more about the Lyapunov spectrum. In particular, by the main theorem of Eskin-Matheus, we conclude simplicity of the Lyapunov spectrum. That is, the spectrum is of the form

$$1 > \lambda_2 > \lambda_3 > \lambda_4 > 0 > -\lambda_4 > -\lambda_3 > -\lambda_2 > -1.$$

Some numerical experiments with the `surface_dynamics` package [2] indicate that

$$\lambda_2 \sim 0.531505241984150,$$

$$\lambda_3 \sim 0.277626450819842,$$

$$\lambda_4 \sim 0.191386026412578.$$

Remark 5.2. *For arithmeticity, the article [1] used 2-cylinder decompositions of their origami, but for our origami no such decomposition exists. Indeed, there are 4 cylinders in the horizontal direction and any other rational direction can be moved to the horizontal using the action of the Veech group $\mathrm{SL}(2, \mathbb{Z})$. The more relevant data for the metacode of [1] is that the homological dimension is 2, so that*

the associated Dehn multitwist C will have a specific form. Namely, the image of the restriction of $C - Id$ to the zero holonomy part is a one-dimensional subspace.

Theorem 5.3. *The monodromy group Γ is arithmetic.*

Proof. Using the metacode of [1], it suffices to find 3 transvections with respect to 3 rational directions of cylinder decomposition. Recall, by the argument from Proposition 2.3, that in any rational direction there are two “long” cylinders with homologous waist curves and two “short” cylinders with homologous waist curves. We consider three rational directions indicated by the vectors $(1, 0)$, $(0, 1)$, and $(1, 2)$ and consider linear combinations of the waist curves so that we are in the zero-holonomy subspace. The corresponding zero-holonomy vectors of the rational directions $(1, 0)$, $(0, 1)$, and $(1, 2)$ are $w_1 = 2\gamma_3 - 6\gamma_1$, $w_2 = 2\gamma_7 - 6\gamma_5$, and $w_3 = 2\alpha - 6\beta$.

Recall the basis $\{\epsilon_1, \dots, \epsilon_6\}$ of zero-holonomy subspace from the end of Section 3. With respect to this basis, the curves chosen for the transvection are given by $w_1 = 2\epsilon_1$, $w_2 = 2\epsilon_5 - 6\epsilon_3$, and $w_3 = -6\epsilon_1 + 12\epsilon_3 + 8\epsilon_4 - 4\epsilon_5 + 8\epsilon_6$.

The transvections C_{w_i} in the directions w_i for $i = 1, 2, 3$ act on the zero-holonomy subspace are given by

$$\begin{aligned} C_{w_1}(X) &= X + 2\langle X, \gamma_3 \rangle \gamma_3 + 6\langle X, \gamma_1 \rangle \gamma_1, \\ C_{w_2}(X) &= X + 2\langle X, \gamma_7 \rangle \gamma_7 + 6\langle X, \gamma_5 \rangle \gamma_5, \\ C_{w_3}(X) &= X + 2\langle X, \alpha \rangle \alpha + 6\langle X, \beta \rangle \beta. \end{aligned}$$

Let e be the annihilator of the subspace $W = \text{span}_{\mathbb{Q}}\{w_1, w_2, w_3\}$. It can be computed by the formula,

$$e = -\frac{\langle w_3, w_2 \rangle}{\langle w_1, w_2 \rangle} w_1 - \frac{\langle w_3, w_1 \rangle}{\langle w_2, w_1 \rangle} w_2 + w_3 = -w_1 + 2w_2 + w_3.$$

Then the transvections C_{w_i} for $i = 1, 2, 3$ with respect to the ordered basis $\{w_1, w_3, e\}$ are

$$C_{w_1} = \begin{pmatrix} 1 & -12 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, C_{w_2} = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 2 & 0 \\ -1 & -1 & 1 \end{pmatrix}, C_{w_3} = \begin{pmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

We find a non-trivial element in $\text{Sp}(W)$, by considering the word

$$C_{w_1} C_{w_2} C_{w_1}^{-1} C_{w_2}^{-1} C_{w_3}^3 C_{w_1}^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 12 & 144 & 1 \end{pmatrix}.$$

The metacode of [1] implies that Γ is arithmetic. □

We now explain the process of recording the index and congruence level of Γ . See also Section 6.2 of [1]. Recall that our intersection matrix Ω has determinant 16. Thus, we cannot conjugate Γ to a subgroup of $\text{Sp}(6, \mathbb{Z})$. However, we can pass to a specific finite index subgroup that is conjugate to a subgroup of $\text{Sp}(6, \mathbb{Z})$. These steps allow us to use the software of Detinko–Flannery–Hulpke

[3] and Kattler–Weitze–Schmithüsen [12]. We start this process now by changing our basis Ω to a more standard form. Let

$$\Theta = \begin{pmatrix} 0 & -1 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ -1 & 4 & 0 & 2 & 0 & 0 \end{pmatrix} \text{ so that } \Theta^t \Omega \Theta = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -4 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \end{pmatrix}.$$

That is, Θ is a change of basis into a more standard form with respect to a basis $\{b_1, b_2, b_3, b_4, b_5, b_6\}$. Let $\Gamma' = \Theta^{-1}\Gamma\Theta$. By passing to the sublattice with basis $\{4b_1, b_2, 4b_3, b_4, b_5, b_6\}$ and using the software of Kattler–Weitze–Schmithüsen [12], we obtain an index 288 subgroup of Γ' that can be conjugated to the standard symplectic group. Applying the software of Detinko–Flannery–Hulpke [3] yields the congruence level 64 and the index 4156153952993280 in $\text{Sp}(6, \mathbb{Z})$.

References

- [1] E. BONNAFOUX, M. KANY, P. KATTLER, C. MATHEUS, R. NIÑO, M. SEDANO-MENDOZA, F. VALDEZ, AND G. WEITZE-SCHMITHÜSEN. Arithmeticity of the Kontsevich–Zorich monodromies of certain families of square-tiled surfaces. arXiv:math/220606595. 1414, 1416, 1420, 1421, 1422
- [2] V. DELECROIX, C. FOUGERON, AND S. LELIÈVRE. Surface dynamics - sagemath package, version 0.4.1, 2019. 1421
- [3] A. DETINKO, D. L. FLANNERY, AND A. HULPKE. Zariski density and computing in arithmetic groups. *Math. Comp.*, **87**(310):967–986, 2018. MR3739225, Zbl 1432.20035, arXiv:1611.05921, doi: 10.1090/mcom/3236. 1415, 1420, 1423
- [4] G. FORNI. A geometric criterion for the nonuniform hyperbolicity of the Kontsevich–Zorich cocycle. *J. Mod. Dyn.*, **5**(2):355–395, 2011. With an appendix by Carlos Matheus. MR2820565, Zbl 1238.30031, arXiv:10094655, doi: 10.3934/jmd.2011.5.355. 1417
- [5] G. FORNI AND C. MATHEUS SANTOS. An example of a Teichmüller disk in genus 4 with degenerate Kontsevich–Zorich spectrum arXiv:08100023. 1415
- [6] R. GUTIÉRREZ-ROMO, D. LEE, AND A. SANCHEZ. Kontsevich–Zorich monodromy groups of translation covers of some platonic solids. To appear in *Groups, Geometry, and Dynamics* arXiv:2208.08460. 1419
- [7] F. HERRLICH. Teichmüller curves defined by characteristic origamis. In *The geometry of Riemann surfaces and abelian varieties*, volume 397 of *Contemp. Math.*, pages 133–144. Amer. Math. Soc., Providence, RI, 2006. MR2218004, Zbl 1098.14019 doi: 10.1090/conm/397/07468. 1416, 1417
- [8] F. HERRLICH AND G. SCHMITHÜSEN. An extraordinary origami curve. *Math. Nachr.*, **281**(2):219–237, 2008. MR2387362, Zbl 1159.14012 arXiv:0509195 doi: 10.1002/mana.200510597 1415
- [9] P. HUBERT AND C. MATHEUS SANTOS. An origami of genus 3 with arithmetic Kontsevich–Zorich monodromy. *Math. Proc. Cambridge Philos. Soc.*, **169**(1):19–30, 2020. MR4120783, Zbl 1479.37033, arXiv:180609855, doi: 10.1017/s0305004119000057. 1414, 1415
- [10] M. KANY AND C. MATHEUS. Arithmeticity of the Kontsevich–Zorich monodromies of certain families of square-tiled surfaces II. arXiv:2301.06894. 1414, 1415, 1420, 1421
- [11] P. KATTLER. Index of the Kontsevich–Zorich monodromy of origamis in $\mathcal{H}(2)$. arXiv:2307.00816 1414

- [12] P. KATTLER AND G. WEITZE-SCHMITHÜSEN. Extension of the origami-package: Actions on the homology. *unpublished GAP package*. 1415, 1416, 1420, 1423
- [13] C. MATHEUS. Three lectures on square-tiled surfaces. In *Teichmüller theory and dynamics, Panor. Synthèses*, **58**, 77–99. Soc. Math. France, Paris, (2022). MR4563316, Zbl 07754057. 1415, 1419
- [14] C. MATHEUS AND J.-C. YOCCOZ. The action of the affine diffeomorphisms on the relative homology group of certain exceptionally symmetric origamis. *J. Mod. Dyn.*, **4**(3):453–486, 2010. MR2729331, Zbl 1220.37004, arXiv:09121425, doi: 10.3934/jmd.2010.4.453. 1415
- [15] P. SARNAK. Notes on thin matrix groups. In *Thin groups and superstrong approximation, Math. Sci. Res. Inst. Publ.*, **61**, 343–362. Cambridge Univ. Press, Cambridge, 2014. MR3220897, Zbl 1365.11039, arXiv:1212.3525. 1413
- [16] S. T. SHRESTHA AND J. WANG. Statistics of square-tiled surfaces: symmetry and short loops. *Exp. Math.*, **31**(4):1314–1331, 2022. MR57K20 (30F99 37D40), Zbl 1502.32001, arXiv:1902.08214, doi: 10.1080/10586458.2020.1786864. 1415
- [17] S. SINGH AND T. N. VENKATARAMANA. Arithmeticity of certain symplectic hypergeometric groups. *Duke Math. J.*, **163**(3):591–617, 2014. MR3165424, Zbl 1287.22005, arXiv:1208.6460, doi: 10.1215/00127094-2410655. 1416, 1420

(Xun Gong) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT SAN DIEGO,
9500 GILMAN DR, LA JOLLA, CA 92093, USA
x1gong@ucsd.edu

(Anthony Sanchez) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT SAN DIEGO,
9500 GILMAN DR, LA JOLLA, CA 92093, USA
ans032@ucsd.edu

This paper is available via <http://nyjm.albany.edu/j/2023/29-55.html>.