

# Package ‘rateratio.test’

October 14, 2022

**Type** Package

**Title** Exact Rate Ratio Test

**Version** 1.1

**Date** 2022-05-09

**Author** Michael Fay <mfay@niaid.nih.gov>

**Maintainer** Michael Fay <mfay@niaid.nih.gov>

**Depends** R (>= 2.4.1), stats

**Description** Performs exact rate ratio tests.

**License** GPL-3

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2022-05-09 14:50:02 UTC

## R topics documented:

|                          |          |
|--------------------------|----------|
| rateratio.test . . . . . | 1        |
| <b>Index</b>             | <b>4</b> |

---

rateratio.test            *An Exact Rate Ratio Test Assuming Poisson Counts*

---

## Description

Performs the uniform most powerful unbiased test on the ratio of rates of two Poisson counts with given time (e.g., perons-years) at risk for each count.

## Usage

```
rateratio.test(x, n, RR = 1,  
              alternative = c("two.sided", "less", "greater"),  
              conf.level = 0.95)
```

**Arguments**

|             |   |
|-------------|---|
| x           | a vector of length 2 with counts for the two rates  |
| n           | a vector of length 2 with time at risk in each rate   |
| RR          | the null rate ratio (two.sided) or the rate ratio on boundary between null and alternative  |
| alternative | a character string specifying the alternative hypothesis, must be one of "two.sided" (default), "greater" or "less". You can specify just the initial letter. |
| conf.level  | confidence level of the returned confidence interval. Must be a single number between 0 and 1.  |

**Details**

The `rateratio.test` tests whether the ratio of the first rate (estimated by  $x[1]/n[1]$ ) over the second rate (estimated by  $x[2]/n[2]$ ) is either equal to, less, or greater than RR. Exact confidence intervals come directly from `binom.test`. The two-sided p-value is defined as either 1 or twice the minimum of the one-sided p-values. See Lehmann (1986, p. 152) or `vignette("rateratio.test")`.

For full discussion of the p-value and confidence interval consistency of inferences, see Fay (2010) and `exactci` package.

**Value**

An object of class 'htest' containing the following components:

|             |  |
|-------------|--|
| p.value     | the p-value of the test  |
| estimate    | a vector with the rate ratio and the two individual rates                                  |
| null.value  | the null rate ratio (two.sided) or the rate ratio on boundary between null and alternative |
| conf.int    | confidence interval  |
| alternative | type of alternative hypothesis   |
| method      | description of method  |
| data.name   | description of data  |

**Note**

Much of the error checking code was taken from `prop.test`.

**Author(s)**

Michael Fay

**References**

- Fay, M. P. (2010). Two-sided exact tests and matching confidence intervals for discrete data. *R Journal*, 2(1), 53-58.
- Lehmann, E.L. (1986). *Testing Statistical Hypotheses* (second edition). Wadsworth and Brooks/Cole, Pacific Grove, California.

### See Also

See [poisson.exact](#) in the [exactci](#) package, which gives the same test.

### Examples

```
### p values and confidence intervals are defined the same way
### so there is consistency in inferences
rateratio.test(c(2,9),c(17877,16660))
### Small counts and large time values will give results similar to Fisher's exact test
### since in that case the rate ratio is approximately equal to the odds ratio
### However, for the Fisher's exact test, the two-sided p-value is defined differently from
### the way the confidence intervals are defined and may imply different inferences
### i.e., p-value may say reject OR=1, but confidence interval says not to reject OR=1
fisher.test(matrix(c(2,9,17877-2,16660-9),2,2))
```

# Index

\* **htest**

rateratio.test, 1

exactci, 3

poisson.exact, 3

rateratio.test, 1